

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 078 985 A2**

(12)

## EUROPEAN PATENT APPLICATION

(43) Date of publication:

28.02.2001 Bulletin 2001/09

(51) Int Cl.7: **C12N 15/11**, C12N 9/12,  
C12N 15/63, C12N 15/87

(21) Application number: **00307362.4**

(22) Date of filing: **25.08.2000**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **27.08.1999 US 384162**

(71) Applicant: **Her Majesty in Right of Canada,  
represented by The Minister of Agriculture and  
Agri-Food Canada  
Ottawa, Ontario K1A 0C6 (CA)**

(72) Inventors:

- **Xing, Ti**  
Ottawa, Ontario K2B 6S6 (CA)
- **Malik, Kamal**  
Ottawa, Ontario K1S 5A5 (CA)
- **Martin-Heller, Teresa**  
Gloucester, Ontario K1J 6X2 (CA)
- **Miki, Brian L**  
Ottawa, Ontario K1H 5V1 (CA)

(74) Representative: **Maschio, Antonio  
D Young & Co,  
21 New Fetter Lane  
London EC4A 1DA (GB)**

(54) **Map kinase kinases (MEK)**

(57) A mitogen-activated protein (MAP) kinase kinase gene, tMEK2, was isolated from tomato cv. Bonny Best. By mutagenesis, a permanently-active variant, tMEK2<sup>MUT</sup>, was created. Both wild type tMEK2 and mutant tMEK2<sup>MUT</sup> were driven by a strong constitutive promoter, tCUPΔ, in a tomato protoplast transient expression system. Pathogenesis-related genes, PR1b1 and PR3, and a wound-inducible gene, ER5, were activated by tMEK2<sup>MUT</sup> expression revealing the convergence of

the signal transduction pathways for pathogen attack and mechanical stress at the level of MAPKK. Activation of biotic and abiotic stress response genes downstream of tMEK2 occurred through divergent pathways involving at least two classes of mitogen-activated protein kinase. This study shows that tMEK2 may play an important role in the interaction of signal transduction pathways that mediate responses to both biotic (eg disease) and abiotic (wound responsiveness) stresses.

**EP 1 078 985 A2**

## Description

[0001] The present invention relates to a derivative of a mitogen-activated protein (MAP) kinase kinase and the use of said derivative for increasing disease resistance and enhanced stress tolerance in plants.

## BACKGROUND OF THE INVENTION

[0002] Signaling mechanisms that mediate plant defense responses may be strongly conserved among plants. This is supported by the observation that several classes of R genes confer disease resistance when expressed in heterologous plant species. For instance, the tomato disease resistance gene, *Cf-9*, was shown to confer responsiveness to the fungal avirulence gene product *Avr9* in transgenic tobacco and potato (Hammond-Kosack *et al.*, 1998). Although *Cladosporium fulvum* is exclusively a fungal pathogen of tomato, a rapid hypersensitive response (HR) was induced in transgenic tobacco and potato by experimentally allowing these specific interactions to occur which then induced signaling pathways that could be common to the plants. Furthermore, the tomato disease resistance gene, *Pto*, which specifies race-specific resistance to the bacterial pathogen *Pseudomonas syringae* pv *tomato* carrying the *avrPto* gene, also increased the resistance of tomato to *Xanthomonas campestris* pv *vesicatoria* and *Cladosporium fulvum* when over expressed (Tang *et al.*, 1999). Clearly, it is the recognition of the pathogen that is unique to most plant species; whereas, the defense response is similar among them.

[0003] Considerable progress has now been made in understanding the signal transduction pathways following perception of biotic and abiotic stresses and the information is being used to develop strategies for modifying transgenic plants. Separate manipulations of the G protein pathway (Xing *et al.*, 1996, 1997) may elevate pathogen resistance or induce defense reactions in transgenic tobacco (Beffa *et al.*, 1995) and increase resistance to tobacco mosaic virus infection (Sano *et al.*, 1994). Multiple roles for MAPK (mitogen-activated protein kinase) in plant signal transduction have also been shown, including responsiveness to pathogens, wounding and other abiotic stresses, as well as plant hormones such as ABA, auxin and ethylene (Hirt, 1997; Kovtun *et al.*, 1998). MAPKK (mitogen-activated protein kinase kinase) from *Arabidopsis* (AtMEK1) and tomato (LeMEK1) have been shown to be induced by wounding (Morris *et al.*, 1997; Hackett *et al.*, 1998), and the maize (ZmMEK1) gene was induced by high salinity and cold (Hardin and Wolniak, 1998). These enzymes interact within MAP kinase pathways that are extensively used for transcytoplasmic signaling to the nucleus. In the MAPK signal transduction cascade, MAPKK (MAP kinase kinase) is activated by upstream MAP-KKK (mitogen-activated protein kinase kinase kinase) and in turn activates MAPK. The transcription of specific genes is induced by MAPK through phosphorylation and activation of transcription factors. This pathway has not yet been manipulated in plants.

## SUMMARY OF THE INVENTION

[0004] The present invention relates to a derivative of a mitogen-activated protein (MAP) kinase kinase and the use of said derivative for increasing disease resistance and enhanced stress tolerance in plants.

[0005] According to the present invention it was determined that mutagenesis of a core phosphorylation site of a member of the MAPK cascade can create a permanently-active form, which stimulates both pathogen- and wound-inducible genes when introduced into plant cells.

[0006] Thus, according to the present invention there is provided a nucleic acid sequence encoding a derivative of a mitogen-activated protein kinase kinase gene from plants, wherein said derivative contains a negative charge in a core phosphorylation site of said protein kinase kinase gene.

[0007] Further according to the present invention there is provided a derivative of a mitogen-activated protein kinase kinase gene from plants, wherein said derivative contains a negative charge in a core phosphorylation site of said protein kinase kinase gene.

[0008] In a further embodiment of the present invention there is provided a cloning vector comprising a nucleic acid sequence encoding a derivative of a mitogen-activated protein kinase kinase gene from plants, wherein said derivative contains a negative charge in a core phosphorylation site of said protein kinase kinase gene.

[0009] The present invention also includes a transgenic plant comprising a nucleic acid sequence encoding a derivative of a mitogen-activated protein kinase kinase gene, wherein said derivative contains a negative charge in a core phosphorylation site of said protein kinase kinase gene.

[0010] Further, according to the present invention there is provided a method of increasing disease resistance or enhancing stress tolerance in a plant by introducing into said plant a nucleic acid sequence encoding a derivative of a mitogen-activated protein kinase kinase gene, wherein said derivative contains a negative charge in a core phosphorylation site of said protein kinase kinase gene.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0011] These and other features of the invention will become more apparent from the following description in which reference is made to the appended drawings wherein:

[0012] **FIGURE 1** shows sequence analysis of tMEK2. **FIGURE 1a** shows the DNA (SEQ ID NO:1) and deduced amino acid sequence (SEQ ID NO: 2). Roman numerals under the sequence indicate the 11 subdomains found in protein kinases. The asterisk indicates stop codon. **FIGURE 1b** shows the alignment of the deduced amino acid sequences from catalytic domains of MAPKK subfamily members (SEQ ID NO: 3 to 21). **FIGURE 1c** shows the alignment of amino acid sequences of tMEK2 with other MAPKKs between subdomain VII and VIII. Dashes represent gaps introduced for maximum matching. The amino acid residues in bold and italics between subdomain VII and VIII show putative phosphorylation sites.

[0013] **FIGURE 2** shows the autophosphorylation and substrate phosphorylation activity of tMEK2. **FIGURE 2a** shows the autophosphorylation of tMEK2<sup>WT</sup> and tMEK2<sup>MUT</sup>. Recombinant GST-tMEK2<sup>WT</sup> or GST-tMEK2<sup>MUT</sup> proteins were incubated *in vitro* without any protein kinase substrate followed by SDS-PAGE and autoradiography. **FIGURE 2b** shows the phosphorylation of myelin basic protein (MBP) by tMEK2<sup>WT</sup> and tMEK2<sup>MUT</sup>. Recombinant GST-tMEK2<sup>WT</sup> or GST-tMEK2<sup>MUT</sup> proteins were incubated *in vitro* with MBP followed by SDS-PAGE and transfer to nitrocellulose. The upper panel is the autoradiography of the nitrocellulose filter. The lower panel is the immunoblot with anti-GST antibody.

[0014] **FIGURE 3** shows the constructs of tMEK2<sup>WT</sup> or tMEK2<sup>MUT</sup> driven by the constitutive promoter tCUPA or control plasmid showing GUS gene driven by the constitutive promoter tCUPA.

[0015] **FIGURE 4** shows the expression of tMEK2 in tomato leaf mesophyll protoplasts. The effect was analysed by quantitative RT-PCR following transient expression of tMEK2 in protoplasts. C1, no electroporation; C2, electroporation of control plasmid; MEK2<sup>WT</sup>, electroporation of plasmid with tMEK2<sup>WT</sup> driven by the tCUPA promoter, electroporation of plasmid with tMEK2<sup>MUT</sup> driven by tCUPA promoter. The pathogenesis-related genes PR1b1, PR3 and Twil were tested. Tomato actin was used as an internal standard.

[0016] **FIGURE 5** shows the activation of ER5 by tMEK2. **FIGURE 5a** shows RNA gel blot analysis of total RNA (15 µg) from leaves following wounding for the indicated time in hours, showing wound-induced activation of tMEK2 and ER5 genes. **FIGURE 5b** shows the activation of ER5 gene by tMEK2. The effect was analysed by quantitative RT-PCR following transient expression of tMEK2 in protoplasts. Lane settings are as described in Figure 4. Tomato actin was used as an internal standard.

[0017] **FIGURE 6** shows the effect of MAPK inhibitors on tMEK2<sup>MUT</sup>-induced gene activation. Kinase inhibitors at the concentration of 1 µM for staurosporine, 350 nM for SB 202190 and 1 µM for PD 98059, SB 203580 and SB 202474 were included in the proteoplast incubation buffer.

[0018] **FIGURE 7** shows the comparison of disease symptoms on a leaf from a wild type plant and on a leaf from tMEK2<sup>MUT</sup> transformed plant.

## DESCRIPTION OF PREFERRED EMBODIMENT

[0019] According to the present invention there is provided a derivative of a mitogen-activated protein kinase kinase (MAPKK). The present invention also relates to a method for increasing disease resistance and enhanced stress tolerance in plants using said derivative.

[0020] When used herein the term derivative means a modified MAPKK protein, wherein said modification includes the replacement of one or more amino acids of the wild type MAPKK with one or more other amino acids. Therefore said derivative is a non-naturally occurring variant of the wild type MAPKK.

[0021] MAPK signaling cascades are ubiquitous among eukaryotes from yeast to human (Guan, 1994) and mediate a large array of signal transduction pathways in plants (Hirt, 1997; Mizoguchi *et al.*, 1997). The cascades utilize the reversible phosphorylation of regulatory proteins to achieve rapid biochemical responses to changing external and internal stimuli. A specific MAPK is rapidly activated by pathways responding to cold, drought, mechanical stimuli and wounding (Bogre *et al.*, 1997; Jonak *et al.*, 1996; Seo *et al.*, 1995; Usami *et al.*, 1995). MAPKs are also rapidly activated by pathways responding to pathogen elicitors (Ligterink *et al.*, 1997; Suzuki and Shinshi, 1995). Other factors such as salicylic acid which is a signaling molecule in the pathogen response, may intervene in the signal cascade by transiently activating a MAPK in tobacco cells (Zhang and Klessig, 1997). MAPKK, which activates MAPK by phosphorylation in the signal cascade has been identified in Arabidopsis, tobacco, maize and tomato (Mizoguchi *et al.*, 1997; Shibata *et al.*, 1995; Hardin and Wolniak, 1998). Although phosphorylation of MAPKK by MAPKKK is the primary mechanism for initiating the signal cascade, regulation at the level of gene expression has also been implied. For instance, transcriptional activity of an Arabidopsis MAPKK, MEK1 (Morris *et al.*, 1997), and a tomato MAPKK, tMEK1 (Hackett *et al.*, 1998), was increased by wounding. Transcriptional activity of ZmMEK1, a maize MAPKK, was slightly increased in roots by high salinity and was substantially decreased by cold (Hardin and Wolniak, 1998). In this study, tomato tMEK2 mRNA accumulation was also induced by wounding of leaves but transient expression in protoplasts did not result in

the activation of the target gene ER5. This observation supported the view that biochemical activation of MAPKK by phosphorylation was the primary factor in signal transduction and that transcriptional control plays a secondary role.

[0022] Yeast and animal MAPKK are activated when serine and serine/threonine residues in the SxAXS/T motif, located upstream of the subdomain VIII are phosphorylated by MAPKKK. The putative consensus motif for characterised plant MAPKK is a S/TxXXxxS/T signature. This motif contains two additional residues when compared with the motif SxAXS/T detected in other eukaryotes. Thus, according to the present invention the use of a plant gene encoding the MAPKK is preferred to that of the yeast and animal genes, as the plant gene provides additional sites for manipulation. The plant genes also provide additional combinations of sites that can be modified according to the present invention. Thus, according to the present invention one or multiple sites of the plant gene can be modified.

[0023] According to the present invention, by creating a negative charge around a core phosphorylation site the activation by MAPKKK was not needed for MAPKK activity.

[0024] According to the present invention possible core phosphorylation sites include: serine and/or threonine sites located upstream of the subdomain VIII.

[0025] According to the present invention the creation of a negative charge around one of said core phosphorylation sites includes replacement of one or more amino acids with an amino acid selected from the group consisting of: any negatively charged amino acids. In one embodiment of the present invention said negatively charged amino acids include glutamic acid and aspartic acid.

[0026] In one embodiment of the present invention MAPKK gene, from various sources can be modified, as described above. As noted earlier MAPK signalling cascades are ubiquitous among eukaryotes from yeast to human. Suitable examples of a suitable gene that can be used according to the present invention include *Lycopersicum esculentum* cv Bonny Best, tMEK2, together with other genes available in the art, as exemplified by the following:

*Arabidopsis thaliana*, AtMAP2K $\alpha$ , (Jouannic S., Hamal A., Kreis M., Henry Y. 1996, Molecular cloning of an *Arabidopsis thaliana* MAP kinase kinase-related cDNA. Plant Physiol. 112:1397)

*A. thaliana*, AtMKK4, (Genbank accession number AB015315)

*A. thaliana*, AtMEK1, (Morris P.C., Cuerrier D., Leung L., Giraudat J. 1997, Cloning and characterisation of MEK1, an *Arabidopsis* gene encoding a homologue of MAP kinase kinase. Plant Mol. Biol. 35: 1057-1064)

*L. esculentum* tomato c.v. Alisa Craig, LeMEK1, (Genbank accession number AJ000728)

*Zea mais*, ZmMEK1, (Genbank accession number U83625)

*A. thaliana*, AtMAP2K $\beta$ , (Genbank accession number AJ006871)

*N. tabacum*, NPK2, (Shibata W., Banno H., Hirano YIK., Irie K. Machida SUC., Machida Y. 1995, A tobacco protein kinase, NPK2, has a domain homologous to a domain found in activators of mitogen-activated protein kinases (MAPKKs). Mol. Gen. Genet. 246: 401-410)

*A. thaliana*, AtMKK3, (Genbank accession number AB015314)

*D. discoideum*, DdMEK1, (Nakai K., Kanehisa M. 1992, A knowledge base for predicting protein localisation sites in eukaryotic cells. Genomics 14:897-911.)

*Leishmania donovani*, LPK, (Li S., Wilson ME., Donelson JE. 1996, *Leishmania chagasi*: a gene encoding a protein kinase with a catalytic domain structurally related to MAP kinase kinase. Exp. Parasitol. 82: 87-96.)

*Drosophila melanogaster*, HEP, (Glise B., Bourbon H., Noselli S. Hemipterous encodes a novel *Drosophila* MAP kinase kinase, required for epithelial cell sheet movement. 1995, Cell 83: 451-461.)

*Homo sapiens*, MEK1, (Zheng C., Guan K. 1993, Cloning and characterisation of two distinct human extracellular signal-regulated kinase activator kinases MEK1 and MEK2. J. Biol. Chem. 268: 11435-11439)

*R. norvegicus*, MEK5, (English JM., Vanderbilt CA., Xu S., Marcus S., Cobb MH. 1995, Isolation of MEK5 and differential expression of alternatively spliced forms. J. Biol. Chem. 270: 28897-28902.)

*H. sapiens*, MKK3, (Derijard B., Raingeaud J., Barrett T., Wu IH., Han J., Ulevitch RJ., Davis RJ. 1995, Independent

human MAPkinase signal transduction pathways defined by MEK and MKK isoforms. Science 267:682-685.)

*Saccharomyces cerevisiae*, PBS2, (Boguslawski G., Polazzi JO. 1987, Complete nucleotide sequence of a gene conferring polymyxin B resistance on yeast: similarity of the predicted polypeptide to protein kinases. Proc. Natl. Acad. Sci. USA 84: 5848-5852.)

*S. cerevisiae*, STE7, (Teague MA., Chaleff DT., Errede B. 1986, Nucleotide sequence of the yeast regulatory gene STE7 predicts a protein homologous to protein kinases. Proc. Natl. Acad. Sci. USA 83: 7371-7375.)

*Candida albicans*, FIST 7, (Clark KL., Feldmann PJ. Dignard D. 1995, Constitutive activation of the *Saccharomyces cerevisiae* mating response pathway by a MAP kinase kinase from *Candida albicans*. Mol. Gen. Genet. 249: 609-621.)

*S. cerevisiae*, MKK1, (Irie T., Takase MKS., Lee KS., Levin DE., Araki H., Matsumoto K., Oshima Y. 1993, MKK1 and MKK2, encoding *Saccharomyces cerevisiae* MAP kinase kinase homologues function in the pathway mediated by protein kinase C. Mol. Cell. Biol. 13:3076-3083.)

[0027] In a further embodiment of the present invention putative phosphorylation activation sites are selected from the group consisting of:

*Lycopersicum esculentum* c.v. Bonny Best, tMEK 2: 219serine, 220threonine, 221serine and 226threonine;  
*Arabidopsis thaliana*, AtMAP2K $\alpha$ : 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMKK4: 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMEK1: 219serine, 220threonine, 221serine, 222serine and 226serine;  
*L. esculentum*, LeMEK1: 219serine, 220threonine, 221serine and 226threonine;  
*Zea mays*, ZmMEK1: 219serine, 220serine and 226threonine;  
*A. thaliana*, At MAP2K $\beta$ : 218threonine, 220threonine and 226threonine;  
*N. tabacum*, NPK2: 219serine, 220serine and 226threonine;  
*A. thaliana*, AtMKK3: 220serine and 226threonine;  
*D. discoideum*, DdMEK1, 220threonine, 222serine and 226threonine;  
*Leishmania donovani*, LPK: 220threonine, 224serine, 225serine and 226threonine;  
*Drosophila melanogaster*, HEP: 220serine and 226threonine;  
*Homo sapiens*, MEK1: 220serine and 226serine;  
*R. norvegicus*, MEK5: 220serine and 226threonine;  
*H. sapiens*, MKK3: 220serine and 226threonine;  
*Saccharomyces cerevisiae*, PBS2: 220serine and 226threonine;  
*S. cerevisiae*, STE7: 220serine and 226threonine;  
*Candida albicans*, HST 7: 220serine and 226threonine; and  
*S. cerevisiae*, MKK1: 220serine, 225threonine and 226threonine;

wherein the amino acid numbering system is based on the tomato gene tMEK2.

[0028] In one further embodiment of the present invention, there is provided a derivative of a mitogen-activated protein kinase kinase gene from tomato cv. Bonny Best, wherein the amino acids serine221 and threonine226 have been replaced with aspartic acid.

[0029] Methods of modifying amino acid sequences are well known in the art. In general terms two primers, one for the 3' end and one for the 5' end are used to amplify the coding region. PCR-based site-directed mutagenesis was then done using the procedure as described by Higuchi (1989). Based on the sequence of the PCR product two PCR reactions are used for its mutagenesis. In PCR reaction 1, a primer containing the appropriate base substitution was used together with the 5' primer to amplify the 5' end of the coding region. In PCR reaction 2, a further primer with the appropriate base substitution was used together with the 3' primer to amplify the 3' end of the coding region. Products from both reactions were then purified and combined for 3' extension. The resulting mutant was then amplified with the original 3' and 5' primers.

[0030] The present invention also includes a suitable cloning vector containing the nucleic acid sequence encoding the derivative of the MAPK gene for transforming suitable plant recipients to increase disease resistance and enhance stress tolerance in plants. Suitable cloning vectors include any cloning vectors, Ti plasmid-derived and standard viral vectors well known in the art.

[0031] The cloning vectors can include various regulatory elements well known in the art. For example the cloning vector of the present invention can further comprise a 3' untranslated region. A 3' untranslated region refers to that

portion of a gene comprising a DNA segment that contains a polyadenylation signal and any other regulatory signals capable of effecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by effecting the addition of polyadenylic acid tracks to the 3' end of the mRNA precursor. Polyadenylation signals are commonly recognized by the presence of homology to the canonical form 5' AATAAA-3' although variations are not uncommon.

[0032] Examples of suitable 3' regions are the 3' transcribed non-translated regions containing a polyadenylation signal of *Agrobacterium* tumor inducing (Ti) plasmid genes, such as the nopaline synthase (*Nos* gene) and plant genes such as the soybean storage protein genes and the small subunit of the ribulose-1, 5-bisphosphate carboxylase (ss-RUBISCO) gene.

[0033] The cloning vector of the present invention can also include further enhancers, either translation or transcription enhancers, as may be required. These enhancer regions are well known to persons skilled in the art, and can include the ATG initiation codon and adjacent sequences. The initiation codon must be in phase with the reading frame of the coding sequence to ensure translation of the entire sequence. The translation control signals and initiation codons can be from a variety of origins, both natural and synthetic. Translational initiation regions may be provided from the source of the transcriptional initiation region, or from the structural gene. The sequence can also be derived from the promoter selected to express the gene, and can be specifically modified so as to increase translation of the mRNA.

[0034] To aid in identification of transformed plant cells, the constructs of this invention may be further manipulated to include plant selectable markers. Useful selectable markers include enzymes which provide resistance to chemicals such as an antibiotic such as gentamycin, hygromycin, kanamycin, or herbicides such as phosphorothycin, glyphosate, chlorsulfuram and the like. Similarly, enzymes providing for production of a compound identifiable by colour change such as *GUS* ( $\beta$ -glucuronidase), or luminescence, such as luciferase are useful.

[0035] A promoter, included in the cloning vector of the present invention, can include a constitutive promoter, which will ensure continued expression of the gene. The nucleic acid sequence encoding the derivative of the MAPK gene can also be under the control of an inducible promoter. Said inducible promoter is triggered by an induction response.

[0036] Generally speaking, an inducible promoter is a promoter that is capable of directly or indirectly activating transcription of one or more DNA sequences or genes in response to an inducer. In the absence of an inducer the DNA sequences or genes will not be transcribed. Typically the protein factor, that binds specifically to an inducible promoter to activate transcription, is present in an inactive form which is then directly or indirectly converted to the active form by the inducer. The inducer can be a chemical agent such as a protein, metabolite, growth regulator, herbicide or phenolic compound or a physiological stress imposed directly by heat, cold, salt, or toxic elements or indirectly through the action of a pathogen or disease agent such as a virus. A plant cell containing an inducible promoter may be exposed to an inducer by externally applying the inducer to the cell or plant such as by spraying, watering, heating or similar methods.

[0037] A constitutive promoter directs the expression of a gene throughout the various parts of a plant and continuously throughout plant development. Examples of known constitutive promoters include those derived from the CaMV 35S and *Agrobacterium* Ti plasmid opine synthase gene (Sanders *et al.*, 1987) or ubiquitin (Christensen *et al.*, 1992). Additionally the constitutive promoter described in WO 97/28268 published August 7, 1997.

[0038] Also considered part of this invention are transgenic plants containing the variant of the present invention. Methods of regenerating whole plants from plant cells are known in the art, and the method of obtaining transformed and regenerated plants is not critical to this invention. In general, transformed plant cells are cultured in an appropriate medium, which may contain selective agents such as antibiotics, where selectable markers are used to facilitate identification of transformed plant cells. Once callus forms, shoot formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be used to establish repetitive generations, either from seeds or using vegetative propagation techniques.

[0039] Besides viral cloning vectors, transformation can also be accomplished by particle bombardment using the nucleic acid sequence encoding the derivative of the MAPK gene. Bombardment is a DNA delivery technique using foreign DNA particles delivered to various plant cells, tissues and species using biolistic device such as gun powder-driven biolistic device (Dupont, Wilmington, DE), gas-driven particle delivery system, microtargeting particle accelerator, an air gun apparatus (Daniell, 1997), helium blasting (Pareddy *et al.*, 1997) and instruments based on electric discharge. Transformation can also be achieved by direct uptake of *Agrobacterium* that contained foreign DNA sequence into plants via stomata in the leaves of stem or roots (Clough *et al.*, 1998).

[0040] A further aspect of the present invention is directed to the use of said nucleic acid sequence encoding the derivative of the MAPK gene to increase disease resistance or to enhance stress tolerance in plants. In this aspect of the invention the nucleic acid is introduced into the plant using any of the methods described above.

[0041] Pathogenesis-related (PR) proteins are intra- and extracellular proteins that accumulate in plant tissues or cultured cells after pathogen attack or elicitor treatment (Bowles, 1990). Using PR gene expression as a marker for the plant defence response, both PR1b1 and the chitinase gene were induced by the derivative of the MAPK gene of

the present invention.

[0042] Furthermore, according to the present invention, the transcription of the tomato ER5 gene, ZG (ABA), drought and wounding (Zegzouti *et al.*, 1997) was induced by the derivative of the MAPK gene of the present invention.

[0043] Thus, according to the present invention the derivative of the MAPK gene of the present invention can activate both pathogen- and wound-related genes.

[0044] The use of said nucleic acid sequence encoding the derivative of the MAPK gene can also be used in combination with other methods to increase disease resistance or to enhance stress tolerance in plants. These other methods could include modification of downstream components for example transcription factors and transcriptional activators. The modification of transcription factors was proven to be an effective means to improve plant stress tolerance. Overexpression of a single stress-inducible transcription factor DREB1A isolated from *Arabidopsis* improved plant drought, salt, and freezing tolerance (Masuga *et al.*, 1999). Overexpression of CBF1, an *Arabidopsis* transcriptional activator, enhanced freezing tolerance (Jaglo-Ottosen *et al.*, 1998). There is potential that modification of transcription factors or transcriptional activators downstream of MAPK in our system will enhance disease resistance and stress tolerance.

[0045] In addition there are some parallel pathways that could contribute to increased disease resistance or to enhanced stress tolerance in plants if used in combination with the modified MAPK pathway of the present invention. An example of another parallel pathway would be calcium dependent protein kinase (CDPK) (Sheen, 1996). CDPK has also been shown to act as a key mediator for cold, salt, drought, dark and ABA stresses. In addition CDPK is involved in primary defence response to pathogen attack. Overexpression of either of two different CDPKs (ATCDPK1 and ATCDPK1a) in maize protoplasts active stress signalling (Sheen, 1996). Thus the co-manipulation of the two pathways should further strengthen the defence ability of the plant.

[0046] The present invention is illustrated by the following examples, which are not to be construed as limiting.

## EXAMPLES

### Example 1: Isolation and Modification of tMEK2.

[0047] RNA was extracted with Extact-A-Plant™ RNA Isolation Kit (CloneTech Laboratories, Inc.) from four-week-old tomato leaves. Reverse transcription was as described in Sambrook *et al.* (1989). Cloning was carried out by PCR using Taq DNA polymerase (Life Technologies Inc.). A MAPKK gene, tMEK2, was isolated from tomato cv. Bonny Best by PCR (Figure 1a) using published MAPKK gene sequences of tomato cv. Ailsa Craig and other plant species. It shares a high level of sequence homology with MAPKKs from other species and tomato cultivars (Figure 1b) but compared with MAPKKs from mammals and yeast, tMEK2 and other plant MAPKKs have two more potential core phosphorylation sites between subdomains VII and VIII (Figure 1c).

[0048] Using PCR-assisted, site-directed mutagenesis, amino acids serine221 and threonine226 were replaced with aspartic acid (Figure 1c) creating a negative charge around the core phosphorylation site so that phosphorylation of MAPKK by upstream MAPKKK is no longer necessary for activity. Two primers (5'-end and 3'-end) that span the coding region of tomato cv Ailsa Craig LeMEK1 were used for the amplification of the MAPKK coding sequence in tomato cv Bonney Best. PCR-based site-directed mutagenesis was carried out as described before (Higuchi, 1989). Based on the sequence of the PCR product, two PCR reactions were run for its mutagenesis. In PCR reaction 1, a primer containing the base substitutions (5'GTATGTGCCGACAAA GTCATTGGCCAGTCCATCTGTGCTT-GCTAGTACTGCACTCACAC3'.SEQ ID NO: 22) was used together with 5'-end primer to amplify a 692 bp fragment corresponding to the 5' region of the cloned MAPKK. In PCR reaction 2, a primer containing the base substitutions (5'GTACTAGCAAGCACAGATGGACTGGCCA ATGACTTTGTGGGCACATACAACCTATATGTC3', SEQ ID NO:23) was used together with 3'-end primer to amplify a 429 bp fragment corresponding to the 3' region of the cloned MAPKK. Products from PCR reaction 1 and 2 were then purified and combined for 3' extension. Mutant tMEK2 was amplified with the original 5'-end primer containing *Bam*HI and *Nco*I restriction sites, and 3'-end primer containing *Sall* and *Sma*I restriction sites. The wild type and mutagenized PCR products were purified from an agarose gel using Elu-Quik DNA Purification Kit (Schleicher & Schuell) and ligated into pre-digested pGEM-T Easy vector. The inserts were digested using *Nco*I/*Sma*I and ligated into pTZ19 tCUPA-GUS-nos3'. This derivative of tCUP promoter was created by the following modifications to the original tCUP: mutation of the sequence, 3' deletion of the sequence, nucleotide addition to the sequence, deletion of an upstream out-of-frame ATG methionine initiator codon from the sequence, deletion of the fusion protein encoded by the tobacco genomic DNA from the sequence, addition of restriction sites to the sequence. In detail, exact nucleotide changes are (numbered relative to the tCUP sequence or to the tCUPΔ (sequence as noted): 2084 CATATGA 2090 (*Nde*I recognition site beginning at 2084 underlined) in the tCUP sequence mutated to 2084 CATAGATCT 2092 (*Bgl*II recognition site beginning at 2087 underlined) in the tCUPΔ sequence deleting one restriction site and one upstream out-of-frame ATG methionine initiator codon while adding another restriction site and two nucleotides; 2171 AATACATGG 2179 in the tCUP sequence mutated to 2173 CCACCATGG 2181 in the tCUPΔ sequence

adding a Kozak consensus motif for translational initiation and an NcoI recognition site at 2176 underlined); 2181 to 2224 (relative to tCUP sequence) of tobacco genomic DNA removed from tCUPA (2183 to 2226 relative to tCUPA), deleting the 3' end of the tCUP sequence and the N-terminal fusion peptide encoded by the tobacco genomic DNA. The tCUPA-GUS-nos construct was created by fusion of the tCUPA sequence with a GUS gene and nos terminator having the sequence 2183 CTCTAGAGGAT CCCCGGGTGGTCAGTCCCTT 2213 3' (SEQ ID NO:24) to the GUS ATG at 2214 on the tCUPA sequence (see Figure 3).

#### Example 2: Expression and Phosphorylation Analysis of Recombinant tMEK2

[0049] For in-frame cloning with GST into the *Bam*HI/*Sall* sites in the pGEX-4T-3 vector (Amersham Pharmacia) subcloned PCR products in pGEM-T Easy vector were digested by *Bam*HI/*Sall* and ligated into pGEX-4T-3 cut with the same enzymes. Sequences of cloned products were confirmed by DNA sequencing. The proteins were expressed as glutathione-S-transferase fusions (GST) and purified by glutathione-agarose (Sigma) affinity chromatography essentially as described in manufacturer's protocol. Protein concentration was determined with a Bio-Rad detection system (Bio-Rad).

[0050] Autophosphorylation assay contained 1 µg of GST-tMEK2<sup>WT</sup> or GST-tMEK2<sup>MUT</sup> in 30 mM Hepes (pH 7.5), 5 mM of MgSO<sub>4</sub>, 5 mM of MnSO<sub>4</sub>, and 1 mM CaCl<sub>2</sub>, 10 mM ATP, and 3 µCi γ-<sup>32</sup>P-ATP (specific activity 222 TBq/mmol) in a total volume of 15 µl. The reaction mixture was incubated at 30°C for 45 min and the reaction was stopped by boiling 3 min in SDS sample buffer. As shown in Figure 2a, both wild type and mutant forms of the tMEK2 enzyme showed autophosphorylation activity.

[0051] Substrate phosphorylation assays contained 1 µg of GST-tMEK2<sup>WT</sup> or GST-tMEK2<sup>MUT</sup>, 2 µg of myelin basic protein (MBP, Life Technologies Inc.), 30 mM Hepes (pH 7.5), 5 mM MgSO<sub>4</sub> and 5 mM MnSO<sub>4</sub>. Reactions were carried out at 30°C for 30 min. Phosphorylated products were separated by 10% SDS-PAGE, transferred to nitrocellulose and autoradiographed. Both the wild type and mutant forms of the tMEK2 enzyme phosphorylated myelin basic protein (MPB) *in vitro* (Figure 2b). Protein immunoblotting was performed as described previously (Xing *et al.*, 1996) using antiGST antibody (Amersham Pharmacia) and alkaline phosphatase-conjugated secondary antibody.

#### Example 3: Activation of pathogen- and wound-related genes by tMEK2

[0052] To examine the effects of tMEK2<sup>WT</sup> and tMEK2<sup>MUT</sup> on the activation of pathogenesis-related (PR) or other pathogen-inducible genes a tomato protoplast transient expression system was developed. Chimeric genes, tCUPA-tMEK2<sup>WT</sup>-nos and tCUPA-tMEK2<sup>MUT</sup>-nos, were constructed using the strong constitutive promoter, tCUPA, which was derived from the tCUP promoter as by modification of the mRNA leader sequence described above. After electroporation, transient expression of potential target genes was detected by quantitative RT-PCR. The genes analysed included PR1b1, which is activated by tomato mosaic virus (Tornero *et al.*, 1997); PR3 (chitinase), which is activated during an incompatible *C. fulvum*-tomato interaction (Danhash *et al.*, 1993); and Twi, which is a pathogen- and wound-inducible gene recently identified in tomato (O'Donnell, *et al.*, 1998).

[0053] The following procedures were used.

##### Protoplast isolation and transformation

[0054] Tomato (*Lycopersicon esculentum* cv Bonny Best) were grown at 80% relative humidity in peat soil in growth cabinets programmed for 16 hr days at 25°C and 8 hr nights at 22°C. Light intensity was controlled at 25 pE m<sup>-2</sup> S<sup>-1</sup> emitted from "cool white" fluorescent lamps (Philip Canada, Scarborough, Ontario). The youngest fully expanded leaves were surface sterilized for 5 min in 4% sodium hypochlorite and rinsed three times with sterile water. The lower epidermis was gently rubbed with Carborundum, rinsed with sterile water and leaf fragments of ca. 1 cm<sup>2</sup> were floated with exposed surface facing an enzyme solution containing 0.15% macerozyme R<sub>10</sub> (Yakult Honsha Co., Japan), 0.3% Cellulase "Onozuka" Rio (Yakult Honsha Co., Japan), 0.4 M sucrose in K3 medium (Maliga *et al.*, 1973). After overnight incubation at 30 °C, the enzyme-protoplast mixture was filtered through a 100 µm nylon sieve, centrifuged at 500 g for 5 min. and floated protoplasts were collected and washed twice with W5 medium (Maliga *et al.*, 1973). The protoplasts were kept on ice in W5 medium for 2 hr before transformation.

[0055] The protoplasts were resuspended in electroporation buffer containing 150mM MgCl<sub>2</sub> and 0.4 M mannitol at a density of 1x10<sup>6</sup> protoplasts/ml and co-electroporated with 12-15 g of pTZ19 carrying tMEK2 gene and pJD300 carrying luciferase gene in a total volume of 500 µl as described by Leckie (1994) with some modifications. Electroporation was performed at 200 volts and 100 µF (Gene Pulser II, Bio-Rad). Protoplasts were then allowed to recover on ice in the dark for 10 min followed by centrifugation at 500 g for 5 min. After removal of the supernatant, the protoplast pellet, with about 500 µl of buffer, was supplemented with another 500 µl protoplast incubation buffer. Protoplasts were incubated in the dark at 30°C for 24 hr.



[0056] Kinase inhibitors (CalBiochem, San Diego, CA) at the concentration of 1  $\mu$ M for staurosporine, 350 nM for SB 202190 and 1  $\mu$ M for PD 98059, SB 203580 and SB 202474, when applicable, were included in the protoplast incubation buffer. The inhibitors did not change protoplast viability (data not shown).

#### 5 Luciferase assay

[0057] Luciferase activity in protoplasts co-electroporated with the constructs under study and luciferase DNA as an internal control were determined for evaluation of transformation efficiency. Protoplasts were lysed in 200  $\mu$ L of LUC extraction buffer (100 mM KPO<sub>4</sub>, 1mM EDTA, 10% glycerol, 0.5% Triton X-100 and 7 mM  $\beta$ -mercaptoethanol, pH 7.8).  
 10 After microfuge centrifugation, the supernatant was collected and a 200  $\mu$ L aliquot of LUC assay buffer (25mM Tricine, 15 mM MgCl<sub>2</sub>, 5mM ATP, BSA 1mg/ml, and 5  $\mu$ L  $\beta$ -mercaptoethanol, pH 7.8) was added to each 20  $\mu$ L aliquot followed by 100  $\mu$ L of luciferin (0.5 mM) as substrate. The reaction was assayed in a luminometer as described (Matthews *et al.*, 1995).

#### 15 Quantitative RT-PCR

[0058] RT-PCR was as described above. The number of PCR cycles corresponded to the high end of the range in which a linear increase in products could be detected (generally 14-16 cycles were used). Reaction products were separated on 1.0 % agarose gels. Southern blot analysis was used to estimate levels of specific amplified products.  
 20 Equivalence of cDNA in different samples was verified using PCR reactions for actin. Primers were designed for PCR according to published sequences for tomato PR-1b1, chitinase, Twi1, ER5 and actin (Tornerio *et al.*, 1997; Danhash *et al.*, 1993; O'Donnell *et al.*, 1998; Zegzouti *et al.* 1997; Moniz de Sa and Drouin, 1996).

[0059] Our results indicated that tomato PR1b1, chitinase and Twi1 genes were activated by tMEK2<sup>MUT</sup>. This indicates that tMEK2 can mediate both pathogen and wound signals. Transient expression of the native tMEK2<sup>WT</sup> gene had no  
 25 effect on the expression of the three target genes (Figure 4), indicating that it is not errantly activated in the protoplast system.

#### Example 4: Induction of the Wound-inducible Gene ER5

[0060] Since MAPK may be the point of convergence of the signal transduction pathways for fungal elicitors and mechanical stress (Romeis *et al.*, 1999) we also examined the induction of the wound-inducible gene, ER5 (Zegzouti *et al.*, 1997). Wounding was carried out by crushing leaves across the lamina and mid-vein using a blunt forceps. RNA was extracted after wounding for the indicated period of time. Fifteen  $\mu$ g of RNA was separated per lane on a denaturing formaldehyde gel. Following transfer to nylon membranes, the blot was hybridized with radio labeled fragment of tMEK2  
 35 coding region or fragment of ER5 coding region. Autoradiography was applied to visualize the hybridization signals (Sambrook *et al.*, 1989).

[0061] Wounding of tomato leaves induced both resident tMEK2 and ER5 genes, mRNA accumulation was detectable in 30 min and lasted for at least 4 hrs (Figure 5a). Transient expression of the mutant tMEK2<sup>MUT</sup> gene in tomato protoplasts also activated ER5 (Figure 5b); however, tMEK2<sup>WT</sup> did not (Figure 5b), showing that elevated transcription of tMEK2 alone was not sufficient for transmitting the wound signal to ER5.  
 40

#### Example 5: Different MAPKs downstream of tMEK2

[0062] To study divergence of the signal pathways downstream of tMEK2 the influence of tMAPK2<sup>MUT</sup> expression in tomato protoplasts was examined in the presence of a broad protein kinase inhibitor (staurosporine) and inhibitors specific to the p38 class MAPK (SB 202190 or SB 203580). Staurosporine inhibited all four genes that were previously activated by tMEK2<sup>MUT</sup>; whereas, inhibitors of p38 class MAPK inhibited the PR3 and ER5 genes but not PR1b1 or Twi1. Furthermore, no effects were observed with SB202474, an inert compound acting as a negative control for MAP kinase inhibition studies, or PD 98059, an inhibitor of the MAP kinase cascade which binds to MAPKKK at a site that  
 50 blocks access to activating enzymes (Alessi *et al.*, 1995). The results, shown in Figure 6, are consistent with the divergence of signal pathway downstream of tMEK2. One of these pathways could include a p38 class MAPK.

#### Example 6: Disease Resistance

[0063] Tomato bacterial pathogen *Pseudomonas syringae* pv *tomato* was infiltrated into tomato leaves and the effect of inoculation was recorded 7 days after inoculation. A representative comparison of disease symptoms on a leaf from a wild-type plant and on a leaf from tMEK2<sup>MUT</sup> transformed plant is shown in Figure 7.

## References

- [0064] Alessi, D.R., Cuenda, A., Cohen, P., Dudley, D.T. and Saltiel, A.R. (1995) PD 098059 is a specific inhibitor of the activation of mitogen-activated protein kinase kinase in vitro and in vivo *J. Biol. Chem.* **270**, 27489-27494.
- 5 [0065] Beffa, R., Szell, M., Menwly, P., Pay, A., Vogeli-Lange, R., Metraux, J.P., Meins, F. and Nagy, F. 1995. Cholera toxin elevates pathogen resistance and induces defense reactions in transgenic tobacco plants. *EMBO Journal* **14**, 5753-5761.
- [0066] Bogre, L., Zwerger, K., Meskiene, I., Binarova, P., Csizmadia, V., Planck, C., Wagner, E., Hirt, H. and Heberle-Bors, E. (1997) The cdc2Ms kinase is differently regulated in the cytoplasm and the nucleus. *Plant Physiol.* **113**, 841-852.
- 10 [0067] Bowles, D.J. (1990) Defense-related proteins in higher plants. *Annu. Rev. Biochem.* **59**, 873-907.
- [0068] Christensen A.H., Sharrock R.A., Quail P.H. 1992, Maize polyubiquitin genes: Structure, thermal perturbation of expression and transcript splicing, and promoter activity following transfer to protoplasts by electroporation. *Plant Mol. Biol.* **18**, 675-689.
- 15 [0069] Clough S.J., Bent A.F. 1998. Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**, 735-743.
- [0070] Danhash, N., Wagemakers, C.A., van Kan, J.A. and De Wit, P.J. (1993) Molecular characterization of four chitinase cDNAs obtained from *cladosporium fulvum*-infected tomato. *Plant Mol. Biol.* **22**, 1017-1029.
- [0071] Daniell H. 1997. Transformation and foreign gene expression in plants mediated by microprojectile bombardment. *Methods in Mol. Biol.* **62** Recombinant Gene Expression Protocols (Tuan R., Ed), Humana Press Inc. Tolowa NJ.
- 20 [0072] Guan, K.L. (1994) The nitrogen activated protein kinase signal transduction pathway: From the cell surface to the nucleus. *Cell. Signal.* **6**, 581-589.
- [0073] Hackett, R.M., Oh, S.A., Morris, P.C. and Grierson, D. (1998) A tomato MAP kinase kinase gene (Accession No AJ000728) differentially regulated during fruit development, leaf senescence and wounding (PGR98-151). *Plant Physiol.* **117**, 1526-1526.
- 25 [0074] Hammond-Kosack, K.E., Tang, S., Harrison, K. and Jones J.D.G. (1998) The tomato *Cf-9* disease resistance gene functions in tobacco and potato to confer responsiveness to the fungal avirulence gene Product *Avr9*. *Plant Cell* **10**, 1251-1266.
- [0075] Hardin, S.C. and Wolniak, S.M. (1998) Molecular cloning and characterization of maize *ZmMEK1*, a protein kinase with a catalytic domain homologous to mitogen- and stress-activated protein kinase kinases. *Planta* **206**, 577-584.
- 30 [0076] Higuchi, R. (1989) Using PCR to engineer DNA. In: *PCR Technology: Principles and Applications for DNA Amplification* (ed. Erlich H.A). Stockton Press, New York.
- [0077] Hirt, H. (1997) Multiple roles of MAP kinases in plant signal transduction. *Trends Plant Sci.* **2**, 11-15.
- [0078] Jaglo-Ottosen K.R., Gilmour S.J., Zarka D.G., Schabenberger O., Thomashow M.F. 1998, *Arabidopsis* CBF1 overexpression induced COR genes and enhances freezing tolerance. *Science* **280**, 104-106.
- 35 [0079] Jonak, C., Kiegl, S., Ligterink, W., Barker, P.J., Huskisson, N.S. and Hirt, H. (1996)
- [0080] Stress signaling in plants: A mitogen-activated protein kinase pathway is activated by cold and drought. *Proc. Natl. Acad. Sci. USA* **93**, 11274-11279.
- [0081] Kasuga M., Liu Q., Miura S., Yamaguchi-Shinozaki K., and Shinozaki K. 1999, Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nature Biotechnology* **17**, 287-291.
- 40 [0082] Kovtun, Y., Chiu, W.-L., Zeng W. and Sheen J. (1998) Suppression of auxin signal transduction by a MAPK cascade in higher plants. *Nature* **395**, 716-720.
- [0083] Ligterink, W., Kroy, T., Nieden, U.Z., Hirt, H. and Scheel D. (1997) Receptor-mediated activation of a MAP kinase in pathogen defense of plants. *Science* **276**, 2054-2057.
- 45 [0084] Maliga, P.S., Breznovitis, A. and Marton, L. (1973) Streptomycin-resistant plants from callus culture of haploid tobacco. *New Biol.* **244**, 29-30.
- [0085] Matthews, B.F., Saunders, J.A., Gelhardt, J.S., Lin, J.J. and Koehle, S.M. (1995) Reporter genes and transient assays for plants. *Methods Mol. Biol.* **55**, 147-162.
- 50 [0086] Mizoguchi, T., Ichimura, K., and Shinozaki, K. (1997) Environmental stress response in plants: the role of mitogen-activated protein kinases. *Trends Biotech.* **15**, 15-19.
- [0087] Moniz de Sa, M., and Drouin, G. (1996) Phylogeny and substitution rates of angiosperm actin genes. *Mol. Biol. Evol.* **13**, 1198-1212.
- [0088] Morris, P.C., Guerrier, D., Leung, J., and Giraudat J. (1997) Cloning and characterisation of *MEK1*, an *Arabidopsis* gene encoding a homologue of MAP kinase kinase. *Plant Mol. Biol.* **35**, 1057-1064.
- 55 [0089] O'Donnell, P.J., Truesdale, M.R., Calvert, C.M., Dorans, A., Roberts, M.R., and Bowles, D.J. (1998) A novel tomato gene that rapidly responds to wound- and pathogen-related signals. *Plant J.* **14**, 137-142.
- [0090] Paredy D., Petolino J., Skokut T., Hopkins N., Miller M., Welter M., Smith K., Clayton D., Pescitelli S.,

Gould A. 1997. Maize transformation via helium blasting. *Maydica* 42, 143-154.

[0091] Romeis, T., Piedras, P., Zhang, S., Klessig, D., Hirt, H., and Jones, J.D.G. (1999) Rapid Avr9- and Cf-9-dependent activation of MAP kinases in tobacco cell cultures and leaves: convergence of resistance gene, elicitor, wound, and salicylate responses. *Plant Cell* 11, 273287.

[0092] Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) Molecular Cloning: a Laboratory Manual, 2nd edn. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press.

[0093] Saunders, P.R., Winter, J.A., Barnason, A.R., Rogers, S.G., Fraley, R.T. (1987), Comparison of cauliflower mosaic virus 35S and nopaline synthase promoters in transgenic plants. *Nucleic Acids Res.* 25, 15, 1543-1558.

[0094] Sano, H., Seo, S., Orudjev, E., Youssefian, S., Ishizuka, K., and Ohashi, Y. (1994) Expression of the gene for a small GTP binding protein in transgenic tobacco elevates endogenous cytokinin levels, abnormally induces salicylic acid in response to wounding and increases resistance to tobacco mosaic virus infection. *Proc. Natl. Acad. Sci.* 91, 1055610560.

[0095] Seo, S., Okamoto, M., Seto, H., Ishizaka, K., Sano, H., and Ohashi, Y. (1995) Tobacco MAP kinase: a possible mediator in wound signal transduction pathways. *Science* 270, 19881992.

[0096] Sheen, J. (1996) Ca<sup>2+</sup>-Dependent Protein Kinases and Stress Signal Transduction in plants. *Science* 274, 1900-1902.

[0097] Shibata, W., Banno H., Ito Y., Hirano K., Irie K., Usami S., Machida C., and Machida, Y. (1995) A tobacco protein kinase, NPK2, has a domain homologous to a domain found in activators of mitogen-activated protein kinases (MAPKKs). *Mol. Gen. Genet.* 246, 401-410.

[0098] Suzuki, K., and Shinshi, H. (1995) Transient activation and tyrosine phosphorylation of a protein kinase in tobacco cells treated with a fungal elicitor. *Plant Cell* 7, 639-647.

[0099] Tang, X., Xie, M., Kin, Y.J., Zhou, J., Klessig, D.F., and Martin, G.B. (1999) Overexpression of Pto activates defense responses and confers broad resistance. *Plant Cell* 11, 15-29.

[0100] Teague, M.A., Chaleff, D.T., and Errede, B. (1986) Nucleotide sequence of the yeast regulatory gene *STE7* predicts a protein homologous to protein kinases. *Proc. Natl. Acad. Sci. USA* 83, 7371-7375.

[0101] Tornero, P., Gadca, J., Coejero, V., and Vera, P. (1997) Two PR-1 genes from tomato are differentially regulated and reveal a novel mode of expression of a pathogenesis-related gene during the hypersensitive response and development. *Mol. Plant-Microbe Interact.* 10, 624634.

[0102] Usami S., Banno H., Ito Y., Nishiha~na R., Machida Y. (1995) Cutting activates a 46-kilodalton protein kinase in plants. *Proc. Natl. Acad. Sci. USA* 92, 8660-8664.

[0103] Xing, T., Higgins, V.J., Blumwald, E. (1996) Regulation of plant defense responses to fungal pathogens: two types of protein kinases in the reversible phosphorylation of the hostplasma membrane H<sup>+</sup>-ATPase. *Plant Cell* 8, 555-564.

[0104] Xing, T., Higgins, V.J., and Blumwald, E. (1997) Identification of G proteins in mediating elicitor-induced dephosphorylation of plasma membrane H<sup>+</sup>-ATPase in host plant. *J. Exp.Bot.* 48, 229-238.

[0105] Zegzouti, H., Jones, B., Marty, C., Lelièvre, J-M., Latché, A., Pech, J-C., and Bonzayen, M. (1997) ER5, a tomato cDNA encoding an ethylene-responsive LEA-like protein: characterization and expression in response to drought, ABA and wounding. *Plant Mol. Biol.* 35, 847-854.

[0106] Zhang, S., and Klessig, D.F. (1997) Salicylic acid activates a 48-kD MAP kinase in tobacco. *Plant Cell* 9, 809-824.

[0107] Zheng, C.F., and Guan, K.L. (1993) Cloning and characterization of two distinct human extracellular signal-regulated kinase activator kinases MEK1 and MEK2. *J. Biol. Chem.* 268, 11435-11439.

[0108] Zheng, C.F. and Guan, K.L. (1994) Activation of MEK family kinases requires phosphorylation of two conserved Ser/Thr residues. *EMBO J* 13, 1123-1131.

[0109] All scientific publications and patent documents are incorporated herein by reference.

[0110] The present invention has been described with regard to preferred embodiments. However, it will be obvious to persons skilled in the art that a number of variations and modifications can be made without departing from the scope of the invention as described in the following claims.

## SEQUENCE LISTING

5

<110> Her Majesty in Right of Canada as Represented by the Minister of  
Agriculture and Agri-Food Canada

10

<120> Novel Plant-Derived Map Kinase Kinase

<130> 08-884280EP

<140>

15

<141>

<150> US 09/384,162

<151> 1999-08-27

20

<160> 24

<170> PatentIn Ver. 2.0

<210> 1

25

<211> 1074

<212> DNA

<213> Lycopersicon esculentum

<220>

30

<221> CDS

<222> (1) .. (1074)

<400> 1

35

atg aag aaa gga tct ttt gca cct aat ctt aaa ctc tct ctt cct cct 48  
Met Lys Lys Gly Ser Phe Ala Pro Asn Leu Lys Leu Ser Leu Pro Pro  
1 5 10 15

40

cct gat gaa gtt gct ctc tcc aaa ttc ctg act gaa tca gga aca ttt 96  
Pro Asp Glu Val Ala Leu Ser Lys Phe Leu Thr Glu Ser Gly Thr Phe  
20 25 30

45

aag gat gga gat ctt ctg gtg aat aga gat gga gtt cga att gtt tcg 144  
Lys Asp Gly Asp Leu Leu Val Asn Arg Asp Gly Val Arg Ile Val Ser  
35 40 45

50

cag agt gaa gtt gca gct cct tca gtt ata cag cca tca gac aac cag 192  
Gln Ser Glu Val Ala Ala Pro Ser Val Ile Gln Pro Ser Asp Asn Gln  
50 55 60

tta tgc tta gct gat ttt gaa gca gta aaa gtt att gga aag gga aat 240  
Leu Cys Leu Ala Asp Phe Glu Ala Val Lys Val Ile Gly Lys Gly Asn  
65 70 75 80

55

EP 1 078 985 A2

	ggt ggt ata gtg cgg ctg gtt cag cat aaa tgg aca ggg caa ttt ttc	288
	Gly Gly Ile Val Arg Leu Val Gln His Lys Trp Thr Gly Gln Phe Phe	
5	85 90 95	
	gct ctc aag gtt att cag atg aat att gat gag tct atg cgc aaa cat	336
	Ala Leu Lys Val Ile Gln Met Asn Ile Asp Glu Ser Met Arg Lys His	
10	100 105 110	
	att gct caa gaa ctg aga att aat cag tca tcc cag tgt cca tat gtt	384
	Ile Ala Gln Glu Leu Arg Ile Asn Gln Ser Ser Gln Cys Pro Tyr Val	
15	115 120 125	
	gtc ata tgc tat cag tcg ttc ttc gac aat ggt gct ata tcc ttg att	432
	Val Ile Cys Tyr Gln Ser Phe Phe Asp Asn Gly Ala Ile Ser Leu Ile	
	130 135 140	
20	ttg gag tat atg gat ggt ggt tcc tta gca gat ttt ctg aaa aag gtc	480
	Leu Glu Tyr Met Asp Gly Gly Ser Leu Ala Asp Phe Leu Lys Lys Val	
	145 150 155 160	
25	aaa aca ata cct gaa cga ttt ctt gct gtt atc tgc aaa cag gtt ctc	528
	Lys Thr Ile Pro Glu Arg Phe Leu Ala Val Ile Cys Lys Gln Val Leu	
	165 170 175	
30	aaa ggc ttg tgg tat ctt cat cat gag aag cat att att cac agg gat	576
	Lys Gly Leu Trp Tyr Leu His His Glu Lys His Ile Ile His Arg Asp	
	180 185 190	
35	ttg aaa cct tcg aat ttg cta atc aat cac aga ggt gat gtc aaa atc	624
	Leu Lys Pro Ser Asn Leu Leu Ile Asn His Arg Gly Asp Val Lys Ile	
	195 200 205	
40	aca gac ttt ggt gtg agt gca gta cta gca agc aca tct gga ctg gcc	672
	Thr Asp Phe Gly Val Ser Ala Val Leu Ala Ser Thr Ser Gly Leu Ala	
	210 215 220	
45	aat acc ttt gtc ggc aca tac aac tat atg tct cca gag aga att tca	720
	Asn Thr Phe Val Gly Thr Tyr Asn Tyr Met Ser Pro Glu Arg Ile Ser	
	225 230 235 240	
50	gga ggt gcc tat gat tac aaa agc gac att tgg agc ttg ggt tta gtc	768
	Gly Gly Ala Tyr Asp Tyr Lys Ser Asp Ile Trp Ser Leu Gly Leu Val	
	245 250 255	
55	ttg ctc gag tgt gca aca ggt cat ttc cca tat aaa cca ccc gag gga	816
	Leu Leu Glu Cys Ala Thr Gly His Phe Pro Tyr Lys Pro Pro Glu Gly	
	260 265 270	

EP 1 078 985 A2

gat gaa gga tgg gtc aat gtc tat gaa ctt atg gaa acc ata gtt gac 864  
 Asp Glu Gly Trp Val Asn Val Tyr Glu Leu Met Glu Thr Ile Val Asp  
 275 280 285  
 5  
 caa cca gaa cct tgt gca cct cct gac caa ttt tct cca caa ttc tgc 912  
 Gln Pro Glu Pro Cys Ala Pro Pro Asp Gln Phe Ser Pro Gln Phe Cys  
 290 295 300  
 10  
 tca ttc ata tct gca tgt gtc cag aag cac cag aag gac aga ctg tcg 960  
 Ser Phe Ile Ser Ala Cys Val Gln Lys His Gln Lys Asp Arg Leu Ser  
 305 310 315 320  
 15  
 gca aat gat ctc atg agt cac cct ttc atc acc atg tac gat gac cag 1008  
 Ala Asn Asp Leu Met Ser His Pro Phe Ile Thr Met Tyr Asp Asp Gln  
 325 330 335  
 20  
 gat atc gat ctt gga tct tac ttc act tcc gca gga cct cca ttg gca 1056  
 Asp ile Asp Leu Gly Ser Tyr Phe Thr Ser Ala Gly Pro Pro Leu Ala  
 340 345 350  
 25  
 aca ctt act gag cta taa 1074  
 Thr Leu Thr Glu Leu  
 355  
 30  
 <210> 2  
 <211> 357  
 <212> PRT  
 <213> Lycopersicon esculentum  
 35  
 <400> 2  
 Met Lys Lys Gly Ser Phe Ala Pro Asn Leu Lys Leu Ser Leu Pro Pro  
 1 5 10 15  
 40  
 Pro Asp Glu Val Ala Leu Ser Lys Phe Leu Thr Glu Ser Gly Thr Phe  
 20 25 30  
 45  
 Lys Asp Gly Asp Leu Leu Val Asn Arg Asp Gly Val Arg Ile Val Ser  
 35 40 45  
 Gln Ser Glu Val Ala Ala Pro Ser Val Ile Gln Pro Ser Asp Asn Gln  
 50 55 60  
 50  
 Leu Cys Leu Ala Asp Phe Glu Ala Val Lys Val Ile Gly Lys Gly Asn  
 65 70 75 80  
 55  
 Gly Gly Ile Val Arg Leu Val Gln His Lys Trp Thr Gly Gln Phe Phe  
 85 90 95

EP 1 078 985 A2

Ala Leu Lys Val Ile Gln Met Asn Ile Asp Glu Ser Met Arg Lys His  
100 105 110

5  
Ile Ala Gln Glu Leu Arg Ile Asn Gln Ser Ser Gln Cys Pro Tyr Val  
115 120 125

10  
Val Ile Cys Tyr Gln Ser Phe Phe Asp Asn Gly Ala Ile Ser Leu Ile  
130 135 140

15  
Leu Glu Tyr Met Asp Gly Gly Ser Leu Ala Asp Phe Leu Lys Lys Val  
145 150 155 160

Lys Thr Ile Pro Glu Arg Phe Leu Ala Val Ile Cys Lys Gln Val Leu  
165 170 175

20  
Lys Gly Leu Trp Tyr Leu His His Glu Lys His Ile Ile His Arg Asp  
180 185 190

25  
Leu Lys Pro Ser Asn Leu Leu Ile Asn His Arg Gly Asp Val Lys Ile  
195 200 205

Thr Asp Phe Gly Val Ser Ala Val Leu Ala Ser Thr Ser Gly Leu Ala  
210 215 220

30  
Asn Thr Phe Val Gly Thr Tyr Asn Tyr Met Ser Pro Glu Arg Ile Ser  
225 230 235 240

35  
Gly Gly Ala Tyr Asp Tyr Lys Ser Asp Ile Trp Ser Leu Gly Leu Val  
245 250 255

Leu Leu Glu Cys Ala Thr Gly His Phe Pro Tyr Lys Pro Pro Glu Gly  
260 265 270

40  
Asp Glu Gly Trp Val Asn Val Tyr Glu Leu Met Glu Thr Ile Val Asp  
275 280 285

45  
Gln Pro Glu Pro Cys Ala Pro Pro Asp Gln Phe Ser Pro Gln Phe Cys  
290 295 300

Ser Phe Ile Ser Ala Cys Val Gln Lys His Gln Lys Asp Arg Leu Ser  
305 310 315 320

50  
Ala Asn Asp Leu Met Ser His Pro Phe Ile Thr Met Tyr Asp Asp Gln  
325 330 335

55  
Asp Ile Asp Leu Gly Ser Tyr Phe Thr Ser Ala Gly Pro Pro Leu Ala  
340 345 350

Thr Leu Thr Glu Leu  
355

5

<210> 3

<211> 225

<212> PRT

10

<213> Arabidopsis thaliana

<400> 3

Leu Asp Met Val Lys Val Ile Gly Lys Gly Ser Ser Gly Val Val Gln  
1 5 10 15

15

Leu Val Gln His Lys Trp Thr Gly Gln Phe Phe Ala Leu Lys Val Ile  
20 25 30

20

Gln Leu Asn Ile Asp Glu Ala Ile Arg Lys Ala Ile Ala Gln Glu Leu  
35 40 45

Lys Ile Asn Gln Ser Ser Gln Cys Pro Asn Leu Val Thr Ser Tyr Gln  
50 55 60

25

Ser Phe Tyr Asp Asn Gly Ala Ile Ser Leu Ile Leu Glu Tyr Met Asp  
65 70 75 80

30

Gly Gly Ser Leu Ala Asp Phe Leu Lys Ser Val Lys Arg His Ile Ile  
85 90 95

His Arg Asp Leu Lys Pro Ser Asn Leu Leu Ile Asn His Arg Gly Glu  
100 105 110

35

Val Lys Ile Thr Asp Phe Gly Val Ser Thr Val Met Thr Asn Thr Ala  
115 120 125

40

Gly Leu Ala Asn Thr Phe Val Gly Thr Tyr Asn Tyr Met Ser Pro Glu  
130 135 140

Arg Ile Val Gly Asn Lys Tyr Gly Asn Lys Ser Asp Ile Trp Ser Leu  
145 150 155 160

45

Gly Leu Val Val Leu Glu Cys Ala Thr Gly Lys Phe Pro Tyr Ala Pro  
165 170 175

50

Pro Asn Gln Glu Glu Thr Trp Thr Ser Val Phe Glu Leu Met Glu Ala  
180 185 190

Ile Val Asp Gln Pro Pro Pro Ala Leu Pro Ser Gly Asn Phe Ser Pro

55



EP 1 078 985 A2

	195	200	205
5	Glu Leu Ser Ser Phe Ile Ser Thr Cys Leu Gln Lys Glu Pro Asn Ser		
	210	215	220
	Arg		
10	225		
	<210> 4		
	<211> 221		
15	<212> PRT		
	<213> Nicotiana tabacum		
	<400> 4		
20	Met Arg Val Phe Gly Ala Ile Gly Ser Gly Ala Ser Ser Val Val Gln		
	1 5 10 15		
	Arg Ala Ile His Ile Pro Thr His Arg Ile Ile Ala Leu Lys Lys Ile		
25	20 25 30		
	Asn Ile Phe Glu Lys Glu Lys Arg Gln Gln Leu Leu Thr Glu Ile Arg		
	35 40 45		
30	Thr Leu Cys Glu Ala Pro Cys Cys Gln Gly Leu Val Glu Phe Tyr Gly		
	50 55 60		
	Ala Phe Tyr Thr Pro Asp Ser Gly Gln Ile Ser Ile Ala Leu Glu Tyr		
35	65 70 75 80		
	Met Asp Gly Gly Ser Leu Ala Asp Ile Ile Lys Val Arg Lys Arg His		
	85 90 95		
40	Leu Val His Arg Asp Ile Lys Pro Ala Asn Leu Leu Val Asn Arg Lys		
	100 105 110		
	Gly Glu Pro Lys Ile Thr Asp Phe Gly Ile Ser Ala Gly Leu Glu Ser		
45	115 120 125		
	Ser Ile Ala Met Cys Ala Thr Phe Val Gly Thr Val Thr Tyr Met Ser		
	130 135 140		
50	Pro Glu Arg Ile Arg Asn Glu Asn Tyr Ser Tyr Pro Ala Asp Ile Trp		
	145 150 155 160		
	Ser Leu Gly Leu Ala Leu Phe Glu Cys Gly Thr Gly Glu Phe Pro Tyr		
55	165 170 175		

EP 1 078 985 A2

Thr Ala Asn Glu Gly Pro Val Asn Leu Met Leu Gln Ile Leu Asp Asp  
180 185 190

5 Pro Ser Pro Ser Leu Ser Gly His Glu Phe Ser Pro Glu Phe Cys Ser  
195 200 205

10 Phe Ile Asp Ala Cys Leu Lys Lys Asn Pro Asp Asp Arg  
210 215 220

<210> 5  
15 <211> 221  
<212> PRT  
<213> Arabidopsis thaliana

<400> 5  
20 Met Arg Val Phe Gly Ala Ile Gly Ser Gly Ala Ser Ser Val Val Gln  
1 5 10 15

25 Arg Ala Ile His Ile Pro Asn His Arg Ile Leu Ala Leu Lys Lys Ile  
20 25 30

30 Asn Ile Phe Glu Arg Glu Lys Arg Gln Gln Leu Leu Thr Glu Ile Arg  
35 40 45

Thr Leu Cys Glu Ala Pro Cys His Glu Gly Leu Val Asp Phe His Gly  
50 55 60

35 Ala Phe Tyr Ser Pro Asp Ser Gly Gln Ile Ser Ile Ala Leu Glu Tyr  
65 70 75 80

Met Asn Gly Gly Ser Leu Ala Asp Ile Leu Lys Val Thr Lys Arg His  
85 90 95

40 Leu Val His Arg Asp Ile Lys Pro Ala Asn Leu Leu Ile Asn His Lys  
100 105 110

45 Gly Glu Pro Lys Ile Thr Asp Phe Gly Ile Ser Ala Gly Leu Glu Asn  
115 120 125

Ser Met Ala Met Cys Ala Thr Phe Val Gly Thr Val Thr Tyr Met Ser  
130 135 140

50 Pro Glu Arg Ile Arg Asn Asp Ser Tyr Ser Tyr Pro Ala Asp Ile Trp  
145 150 155 160

55 Ser Leu Gly Leu Ala Leu Phe Glu Cys Gly Thr Gly Glu Phe Pro Tyr  
165 170 175

**EP 1 078 985 A2**

	Ile	Ala	Asn	Glu	Gly	Pro	Val	Asn	Leu	Met	Leu	Gln	Ile	Leu	Asp	Asp	
5																	
	Pro	Ser	Pro	Thr	Pro	Pro	Lys	Gln	Glu	Phe	Ser	Pro	Glu	Phe	Cys	Ser	
10																	
	Phe	Ile	Asp	Ala	Cys	Leu	Gln	Lys	Asp	Pro	Asp	Ala	Arg				
15																	
	<210>	6															
	<211>	286															
	<212>	PRT															
	<213>	Dictyostelium discoideum															
20																	
	<400>	6															
	Leu	Lys	Ile	Ile	Arg	Val	Leu	Gly	Arg	Gly	Ala	Gly	Gly	Val	Val	Lys	
25																	
	Leu	Ala	Tyr	His	Glu	Thr	Ser	Gly	Thr	Tyr	Ile	Ala	Leu	Lys	Val	Ile	
30																	
	Thr	Leu	Asp	Ile	Gln	Glu	Asn	Ile	Arg	Lys	Gln	Ile	Ile	Leu	Glu	Leu	
35																	
	Lys	Thr	Leu	His	Lys	Thr	Ser	Tyr	Pro	Tyr	Ile	Val	Ser	Phe	Tyr	Asp	
40																	
	Ala	Phe	Tyr	Thr	Glu	Gly	Ser	Ile	Phe	Ile	Ala	Leu	Glu	Phe	Met	Glu	
45																	
	Leu	Gly	Ser	Leu	Ser	Asp	Ile	Met	Lys	Lys	Thr	Ser	Leu	His	Leu	Ile	
50																	
	His	Arg	Asp	Ile	Lys	Pro	Ser	Asn	Ile	Leu	Val	Asn	Asn	Lys	Gly	Glu	
55																	
	Ala	Lys	Ile	Ala	Asp	Phe	Gly	Val	Ser								

EP 1 078 985 A2

Ile Ala Asn Glu Gly Pro Val Asn Leu Met Leu Gln Ile Leu Asp Asp  
180 185 190

5 Pro Ser Pro Thr Pro Pro Lys Gln Glu Phe Ser Pro Glu Phe Cys Ser  
195 200 205

10 Phe Ile Asp Ala Cys Leu Gln Lys Asp Pro Asp Ala Arg  
210 215 220

<210> 6  
15 <211> 286  
<212> PRT  
<213> Dictyostelium discoideum

20 <400> 6  
Leu Lys Ile Ile Arg Val Leu Gly Arg Gly Ala Gly Gly Val Val Lys  
1 5 10 15

25 Leu Ala Tyr His Glu Thr Ser Gly Thr Tyr Ile Ala Leu Lys Val Ile  
20 25 30

Thr Leu Asp Ile Gln Glu Asn Ile Arg Lys Gln Ile Ile Leu Glu Leu  
35 40 45

30 Lys Thr Leu His Lys Thr Ser Tyr Pro Tyr Ile Val Ser Phe Tyr Asp  
50 55 60

35 Ala Phe Tyr Thr Glu Gly Ser Ile Phe Ile Ala Leu Glu Phe Met Glu  
65 70 75 80

Leu Gly Ser Leu Ser Asp Ile Met Lys Lys Thr Ser Leu His Leu Ile  
85 90 95

40 His Arg Asp ile Lys Pro Ser Asn Ile Leu Val Asn Asn Lys Gly Glu  
100 105 110

45 Ala Lys Ile Ala Asp Phe Gly Val Ser Gly Gln Leu Gln His Thr Leu  
115 120 125

Ser Lys Ala Val Thr Trp Val Gly Thr Val Thr Tyr Met Ser Pro Glu  
130 135 140

50 Arg Ile Ser Gly Arg Ser Tyr Ser Phe Asp Ser Asp Ile Trp Ser Leu  
145 150 155 160

55 Gly Leu Thr Ile Leu Glu Cys Ala Ile Gly Lys Phe Pro Tyr Gly Ser

EP 1 078 985 A2

		165		170		175										
5	Asn	Leu	Pro	His	Gln	Gln	Gln	Gln	Pro	Leu	Gln	Gln	Gln	Leu	Gln	Asn
				180					185					190		
	Leu	Asp	Ile	Asn	Asn	Ser	Asn	Asn	Asn	Ile	Arg	Asn	Ser	Asn	Asn	Asn
			195						200					205		
10	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn
			210						215					220		
15	Asn	Asn	Val	Leu	Asp	Ile	Ser	Asn	Gly	Gly	Leu	Val	Asp	Ser	Gly	Ser
	225					230					235					240
	Ser	Val	Pro	Glu	Gly	Met	Gly	Phe	Trp	Val	Leu	Leu	Asp	Cys	Ile	Val
					245					250					255	
20	Lys	Glu	Glu	Val	Pro	Ile	Leu	Pro	Ser	Thr	Phe	Ser	Lys	Glu	Phe	Arg
				260					265					270		
25	Ser	Phe	Ile	Ser	Glu	Cys	Leu	Gln	Lys	Glu	Pro	Thr	Glu	Arg		
			275					280					285			
30	<210>	7														
	<211>	222														
	<212>	PRT														
	<213>	Leishmania donovani														
35	<400>	7														
	Tyr	Ser	Ser	Lys	Arg	Asn	Val	Gly	Ala	Gly	Ala	Ser	Gly	Asp	Val	Phe
	1				5					10				15		
40	Phe	Ala	Arg	Leu	Lys	Asn	Gly	Thr	Ser	Ile	Ala	Leu	Lys	Arg	Ile	Pro
				20					25					30		
	Ile	Ser	Ser	Lys	Ala	His	Arg	Asp	Glu	Val	Asp	Arg	Glu	Leu	Gln	Val
45			35					40					45			
	Phe	Met	Ala	Arg	Ala	Asp	Ser	Pro	Tyr	Val	Met	Asn	Asn	Tyr	Gly	Ala
		50					55					60				
50	Phe	Trp	Asp	Ala	Glu	Asp	Asp	Ala	Ile	Val	Ile	Pro	Met	Glu	Trp	Met
	65					70					75				80	
	Pro	Tyr	Thr	Val	Lys	Asp	Leu	Gly	Leu	Phe	Trp	Gly	Gly	Lys	Arg	Val
55					85					90					95	

**EP 1 078 985 A2**

	Leu	His	Arg	Asp	Leu	Lys	Pro	Ser	Asn	Leu	Leu	Ile	Ser	Glu	Thr	Gly	
5																	
	His	Val	Lys	Ile	Ala	Asp	Phe	Gly	Val	Ser	Lys	Leu	Ile	Gln	Thr	Leu	
10	Ala	Val	Ser	Ser	Thr	Tyr	Val	Ala	Thr	Met	Cys	Phe	Met	Ala	Pro	Glu	
	Arg	Leu	Glu	Gln	Gly	Met	Tyr	Gly	Phe	Ser	Ser	Asp	Val	Trp	Ser	Leu	
15																	
	Gly	Leu	Thr	Met	Ile	Gly	Ala	Val	Thr	Gly	Lys	Asn	Pro	Trp	Ala	Pro	
20	Pro	Glu	Glu	Met	Asn	Leu	Tyr	Gln	Leu	Leu	Gly	Lys	Met	Ala	Asn	Gly	
	Ser	Thr	Pro	Thr	Leu	Pro	Lys	Ser	Gly	Ala	Phe	Ser	Asp	Asp	Val	Lys	
25																	
	Asp	Phe	Val	Lys	Gln	Cys	Leu	Glu	Arg	Asp	Pro	Asp	Thr	Arg			
30																	
	<210>	8															
	<211>	222															
	<212>	PRT															
	<213>	Drosophila melanogaster															
35																	
	<400>	8															
	Leu	Lys	His	Leu	Gly	Asp	Leu	Gly	Asn	Gly	Thr	Ser	Gly	Asn	Val	Val	
40																	
	Lys	Met	Met	His	Leu	Ser	Ser	Asn	Thr	Ile	Ile	Ala	Val	Lys	Gln	Met	
45	Arg	Arg	Thr	Gly	Asn	Ala	Glu	Glu	Asn	Lys	Arg	Ile	Leu	Met	Asp	Leu	
	Asp	Val	Val	Leu	Lys	Ser	His	Asp	Cys	Lys	Tyr	Ile	Val	Lys	Cys	Leu	
50																	
	Gly	Cys	Phe	Val	Arg	Asp	Pro	Asp	Val	Trp	Ile	Cys	Met	Glu	Leu	Met	

EP 1 078 985 A2

His Arg Asp Val Lys Pro Ser Asn Ile Leu Ile Asp Glu Arg Gly Asn  
100 105 110

5 Ile Lys Leu Cys Asp Phe Gly Ile Ser Gly Arg Leu Val Asp Ser Lys  
115 120 125

10 Ala Asn Thr Arg Ala Gly Cys Ala Ala Tyr Met Ala Pro Glu Arg Ile  
130 135 140

Asp Pro Lys Lys Pro Lys Tyr Asp Ile Arg Ala Asp Val Trp Ser Leu  
145 150 155 160

15 Gly Ile Thr Leu Val Glu Leu Ala Thr Ala Arg Ser Pro Tyr Glu Gly  
165 170 175

20 Cys Asn Thr Asp Phe Glu Val Leu Thr Lys Val Leu Asp Ser Glu Pro  
180 185 190

Pro Cys Leu Pro Tyr Gly Glu Gly Tyr Asn Phe Ser Gln Gln Phe Arg  
195 200 205

25 Asp Phe Val Ile Lys Cys Leu Thr Lys Asn His Gln Asp Arg  
210 215 220

30 <210> 9  
<211> 234  
<212> PRT  
35 <213> Homo sapiens

<400> 9  
Phe Glu Lys Ile Ser Glu Leu Gly Ala Gly Asn Gly Gly Val Val Phe  
1 5 10 15

40 Lys Val Ser His Lys Pro Ser Gly Leu Val Met Ala Arg Lys Leu Ile  
20 25 30

45 His Leu Glu Ile Lys Pro Ala Ile Arg Asn Gln Ile Ile Arg Glu Leu  
35 40 45

Gln Val Leu His Glu Cys Asn Ser Pro Tyr Ile Val Gly Phe Tyr Gly  
50 55 60

Ala Phe Tyr Ser Asp Gly Glu Ile Ser Ile Cys Met Glu His Met Asp  
65 70 75 80

55 Gly Gly Ser Leu Asp Gln Val Leu Lys Lys Ala Gly His Lys Ile Met

EP 1 078 985 A2

	85	90	95
5	His Arg Asp Val Lys Pro Ser Asn Ile Leu Val Asn Ser Arg Gly Glu		
	100	105	110
10	Ile Lys Leu Cys Asp Phe Gly Val Ser Gly Gln Leu Ile Asp Ser Met		
	115	120	125
15	Ala Asn Ser Phe Val Gly Thr Arg Ser Tyr Met Ser Pro Glu Arg Leu		
	130	135	140
20	Gln Gly Thr His Tyr Ser Val Gln Ser Asp Ile Trp Ser Met Gly Leu		
	145	150	155
25	Ser Leu Val Glu Met Ala Val Gly Arg Tyr Pro Ile Pro Pro Pro Asp		
	165	170	175
30	Ala Lys Glu Leu Glu Leu Met Phe Gly Gly Met Asp Ser Arg Pro Pro		
	180	185	190
35	Met Ala Ile Phe Glu Leu Leu Asp Tyr Ile Val Asn Glu Pro Pro Pro		
	195	200	205
40	Lys Leu Pro Ser Gly Val Phe Ser Leu Glu Phe Gln Asp Phe Val Asn		
	210	215	220
45	Lys Cys Leu Ile Lys Asn Pro Ala Glu Arg		
	225	230	
50	<210> 10		
	<211> 177		
	<212> PRT		
	<213> Rattus norvegicus		
55	<400> 10		
	Ile Arg Tyr Arg Asp Thr Leu Gly His Gly Asn Gly Gly Thr Val Tyr		
	1	5	10
60	Lys Ala Tyr His Val Pro Ser Gly Lys Ile Leu Ala Val Lys Val Ile		
	20	25	30
65	Leu Leu Asp Ile Thr Leu Glu Leu Gln Lys Gln Ile Met Ser Glu Leu		
	35	40	45
70	Glu Ile Leu Tyr Lys Cys Asp Ser Ser Tyr Ile Ile Gly Phe Tyr Gly		
	50	55	60



EP 1 078 985 A2

Ala Phe Phe Val Glu Asn Arg Ile Ser Ile Cys Thr Glu Phe Met Asp  
65 70 75 80

5 Gly Gly Ser Leu Asp Val Tyr Arg Lys Ile Leu Lys Ile Leu His Arg  
85 90 95

10 Asp Val Lys Pro Ser Asn Met Leu Val Asn Thr Ser Gly Gln Val Lys  
100 105 110

Leu Cys Asp Phe Gly Val Ser Thr Gln Leu Val Asn Ser Ile Ala Lys  
115 120 125

15 Thr Tyr Val Gly Thr Asn Ala Tyr Met Ala Pro Glu Arg Ile Ser Gly  
130 135 140

20 Glu Gln Tyr Gly Ile His Ser Asp Val Trp Ser Leu Gly Ile Ser Phe  
145 150 155 160

Met Glu Leu Ala Leu Gly Arg Phe Pro Tyr Pro Gln Ile Gln Lys Asn  
165 170 175

25 Gln

30 <210> 11  
<211> 185  
<212> PRT  
<213> Homo sapiens

35 <400> 11  
Leu Val Thr Ile Ser Glu Leu Gly Arg Gly Ala Tyr Gly Val Val Glu  
1 5 10 15

40 Lys Val Arg His Ala Gln Ser Gly Thr Ile Met Ala Val Lys Arg Ile  
20 25 30

45 Arg Ala Thr Val Asn Ser Gln Glu Gln Lys Arg Leu Leu Met Asp Leu  
35 40 45

Asp Ile Asn Met Arg Thr Val Asp Cys Phe Tyr Thr Val Thr Phe Tyr  
50 55 60

50 Gly Ala Leu Phe Arg Glu Gly Asp Val Trp Ile Cys Met Glu Leu Met  
65 70 75 80

55 Asp Thr Ser Leu Asp Lys Phe Tyr Arg Lys Val Leu Asp Lys Asn Met  
85 90 95

EP 1 078 985 A2

5 Leu Ser Val Ile His Arg Asp Val Lys Pro Ser Asn Val Leu Ile Asn  
 100 105 110  
 Lys Glu Gly His Val Lys Met Cys Asp Phe Gly Ile Ser Gly Tyr Leu  
 115 120 125  
 10 Val Asp Ser Val Ala Lys Thr Met Asp Ala Gly Cys Lys Pro Tyr Met  
 130 135 140  
 Ala Pro Glu Arg Ile Asn Pro Glu Leu Asn Gln Lys Gly Tyr Asn Val  
 145 150 155 160  
 15 Lys Ser Asp Val Trp Ser Leu Gly Ile Thr Met Ile Glu Met Ala Ile  
 165 170 175  
 20 Leu Arg Phe Pro Tyr Glu Ser Trp Gly  
 180 185  
 25 <210> 12  
 <211> 184  
 <212> PRT  
 <213> *Saccharomyces cerevisiae*  
 30 <400> 12  
 Leu Glu Phe Leu Asp Glu Leu Gly His Gly Asn Tyr Gly Asn Val Ser  
 1 5 10 15  
 35 Lys Val Leu His Lys Pro Thr Asn Val Ile Met Ala Thr Lys Glu Val  
 20 25 30  
 Arg Leu Glu Leu Asp Glu Ala Lys Phe Arg Gln Ile Leu Met Glu Leu  
 35 40 45  
 40 Glu Val Leu His Lys Cys Asn Ser Pro Tyr Ile Val Asp Phe Tyr Gly  
 50 55 60  
 45 Ala Phe Phe Ile Glu Gly Ala Val Tyr Met Cys Met Glu Tyr Met Asp  
 65 70 75 80  
 Gly Gly Ser Leu Asp Lys Ile Tyr Asp Glu Ser Ser Glu Ile Gly His  
 85 90 95  
 50 Asn Ile Ile His Arg Asp Val Lys Pro Thr Asn Ile Leu Cys Ser Ala  
 100 105 110  
 55 Asn Gln Gly Thr Val Lys Leu Cys Asp Phe Gly Val Ser Gly Asn Leu

EP 1 078 985 A2

	115	120	125
5	Val Ala Ser Leu Ala Lys Thr Asn Ile Gly Cys Gln Ser Tyr Met Ala		
	130	135	140
	Pro Glu Arg Ile Lys Ser Leu Asn Pro Asp Arg Ala Thr Tyr Thr Val		
10	145	150	155
	Gln Ser Asp Ile Trp Ser Leu Gly Leu Ser Ile Leu Glu Met Ala Leu		
	165	170	175
15	Gly Arg Tyr Pro Tyr Pro Pro Glu		
	180		
20	<210> 13		
	<211> 189		
	<212> PRT		
	<213> Saccharomyces cerevisiae		
25	<400> 13		
	Leu Val Gln Leu Gly Lys Ile Gly Ala Gly Asn Ser Gly Thr Val Val		
	1 5 10 15		
30	Lys Ala Leu His Val Pro Asp Ser Lys Ile Val Ala Lys Lys Thr Ile		
	20 25 30		
	Pro Val Glu Gln Asn Asn Ser Thr Ile Ile Asn Gln Leu Val Arg Glu		
35	35 40 45		
	Leu Ser Ile Val Lys Asn Val Lys Pro His Glu Asn Ile Ile Thr Phe		
	50 55 60		
40	Tyr Gly Ala Tyr Tyr Asn Gln His Ile Asn Asn Glu Ile Ile Ile Leu		
	65 70 75 80		
	Met Glu Tyr Ser Asp Cys Gly Ser Leu Asp Lys Ile Leu Ser Val Tyr		
45	85 90 95		
	Lys Arg Phe Val Gln Arg Gly Thr Val Tyr Lys Ile Ile His Arg Asp		
	100 105 110		
50	Ile Lys Pro Ser Asn Val Leu Ile Asn Ser Lys Gly Gln Ile Lys Leu		
	115 120 125		
55	Cys Asp Phe Gly Val Ser Lys Lys Leu Ile Asn Ser Ile Ala Asp Thr		
	130 135 140		

EP 1 078 985 A2

Phe Val Gly Thr Ser Thr Tyr Met Ser Pro Glu Arg Ile Gln Gly Asn  
 145 150 155 160  
 5 Val Tyr Ser Ile Lys Gly Asp Val Trp Ser Leu Gly Leu Met Ile Ile  
 165 170 175  
 10 Glu Leu Val Thr Gly Glu Phe Pro Leu Gly Gly His Asn  
 180 185  
 15 <210> 14  
 <211> 189  
 <212> PRT  
 <213> Candida albicans  
 20 <400> 14  
 Leu Leu Thr Leu Lys Gln Leu Gly Ser Gly Asn Ser Gly Ser Val Ser  
 1 5 10 15  
 25 Lys Ile Leu His Ile Pro Thr Gln Lys Thr Met Ala Lys Lys Ile Ile  
 20 25 30  
 His Ile Asp Ser Lys Ser Val Ile Gln Thr Gln Ile Ile Arg Glu Leu  
 35 40 45  
 30 Arg Ile Leu His Glu Cys His Ser Pro Tyr Ile Ile Glu Phe Tyr Gly  
 50 55 60  
 35 Ala Cys Leu Asn Asn Asn Asn Thr Ile Val Ile Cys Met Glu Tyr Cys  
 65 70 75 80  
 Asn Cys Gly Ser Leu Asp Lys Ile Leu Pro Leu Cys Glu Asn His Lys  
 85 90 95  
 40 Ile Ile His Arg Asp Ile Lys Pro Asn Asn Val Leu Met Thr His Lys  
 100 105 110  
 45 Gly Glu Phe Lys Leu Cys Asp Phe Gly Val Ser Arg Glu Leu Thr Asn  
 115 120 125  
 Ser Leu Ala Met Ala Asp Thr Phe Val Gly Thr Ser Met Tyr Met Ser  
 130 135 140  
 50 Pro Glu Arg Ile Gln Gly Leu Asp Tyr Gly Val Lys Ser Asp Val Trp  
 145 150 155 160  
 55 Ser Thr Gly Leu Met Leu Ile Glu Leu Ala Ser Gly Val Pro Val Trp  
 165 170 175

EP 1 078 985 A2

Ser Glu Asp Asp Asn Asn Asn Asp Asp Asp Glu Asp Asp  
180 185

5

<210> 15

<211> 187

<212> PRT

10

<213> *Saccharomyces cerevisiae*

<400> 15

15

Ile Glu Thr Leu Gly Ile Leu Gly Glu Gly Ala Gly Gly Ser Val Ser  
1 5 10 15

Lys Cys Lys Leu Lys Asn Gly Ser Lys Ile Phe Ala Leu Lys Val Ile  
20 25 30

20

Asn Thr Leu Asn Thr Asp Pro Glu Tyr Gln Lys Gln Ile Phe Arg Glu  
35 40 45

25

Leu Gln Phe Asn Arg Ser Phe Gln Ser Glu Tyr Ile Val Arg Tyr Tyr  
50 55 60

30

Gly Met Phe Thr Asp Asp Glu Asn Ser Ser Ile Tyr Ile Ala Met Glu  
65 70 75 80

Tyr Met Gly Gly Arg Ser Leu Asp Ala Ile Tyr Lys Asn Leu Leu Glu  
85 90 95

35

Arg Gly Gly Lys Lys Val Ile His Arg Asp Ile Lys Pro Gln Asn Ile  
100 105 110

40

Leu Leu Asn Glu Asn Gly Gln Val Lys Leu Cys Asp Phe Gly Val Ser  
115 120 125

Gly Glu Ala Val Asn Ser Leu Ala Thr Thr Phe Thr Gly Thr Ser Phe  
130 135 140

45

Tyr Met Ala Pro Glu Arg Ile Gln Gly Gln Pro Tyr Ser Val Thr Ser  
145 150 155 160

50

Asp Val Trp Ser Leu Gly Leu Thr Ile Leu Glu Val Ala Asn Gly Lys  
165 170 175

Phe Pro Cys Ser Ser Glu Lys Met Ala Ala Asn  
180 185

55

EP 1 078 985 A2

<210> 16

<211> 133

<212> PRT

<213> Arabidopsis thaliana

<400> 16

Arg His Ile Val His Arg Asp Ile Lys Pro Ser Asp Leu Leu Ile Asn

1 5 10 15

Ser Ala Lys Asn Val Lys Ile Ala Asp Phe Gly Val Ser Arg Ile Leu

20 25 30

Ala Gln Thr Met Asp Pro Cys Asn Ser Ser Val Gly Thr Ile Ala Tyr

35 40 45

Met Ser Pro Glu Arg Ile Asn Thr Asp Leu Asn His Gly Arg Tyr Asp

50 55 60

Gly Tyr Ala Gly Asp Val Trp Ser Leu Gly Val Ser Ile Leu Glu Phe

65 70 75 80

Tyr Leu Gly Arg Phe Pro Phe Ala Val Ser Arg Gln Gly Asp Trp Ala

85 90 95

Ser Leu Met Cys Ala Ile Cys Met Ser Gln Pro Pro Glu Ala Pro Ala

100 105 110

Thr Ala Ser Gln Glu Phe Arg His Phe Val Ser Cys Cys Leu Gln Ser

115 120 125

Asp Pro Pro Lys Arg

130

<210> 17

<211> 133

<212> PRT

<213> Arabidopsis thaliana

<400> 17

Arg His Ile Val His Arg Asp Ile Lys Pro Ser Asn Leu Leu Ile Asn

1 5 10 15

Ser Ala Lys Asn Val Lys Ile Ala Asp Phe Gly Val Ser Arg Ile Leu

20 25 30

Ala Gln Thr Met Asp Pro Cys Asn Ser Ser Val Gly Thr Ile Ala Tyr

35 40 45

EP 1 078 985 A2

5 Met Ser Pro Glu Arg Ile Asn Thr Asp Leu Asn Gln Gly Lys Tyr Asp  
50 55 60

Gly Tyr Ala Gly Asp Ile Trp Ser Leu Gly Val Ser Ile Leu Glu Phe  
65 70 75 80

10 Tyr Leu Gly Arg Phe Pro Phe Pro Val Ser Arg Gln Gly Asp Trp Ala  
85 90 95

Ser Leu Met Cys Ala Ile Cys Met Ser Gln Pro Pro Glu Ala Pro Ala  
100 105 110

15 Thr Ala Ser Pro Glu Phe Arg His Phe Ile Ser Cys Cys Leu Gln Arg  
115 120 125

20 Glu Pro Gly Lys Arg  
130

25 <210> 18  
<211> 133  
<212> PRT  
<213> Lycopersicon esculentum

30 <400> 18  
Arg Arg Ile Ile His Arg Asp Leu Lys Pro Ser Asn Leu Leu Ile Asn  
1 5 10 15

35 His Arg Gly Glu Val Lys Ile Thr Asp Phe Gly Val Ser Lys Ile Leu  
20 25 30

40 Thr Ser Thr Ser Ser Leu Ala Asn Ser Phe Val Gly Thr Tyr Pro Tyr  
35 40 45

Met Ser Pro Glu Arg Ile Ser Gly Ser Leu Tyr Ser Asn Lys Ser Asp  
50 55 60

45 Ile Trp Ser Leu Gly Leu Val Leu Leu Glu Cys Ala Thr Gly Lys Phe  
65 70 75 80

Pro Tyr Thr Pro Pro Glu His Lys Lys Gly Trp Ser Ser Val Tyr Glu  
85 90 95

50 Leu Val Asp Ala Ile Val Glu Asn Pro Pro Pro Cys Ala Pro Ser Asn  
100 105 110

55 Leu Phe Ser Pro Glu Phe Cys Ser Phe Ile Ser Gln Cys Val Gln Lys

EP 1 078 985 A2

	115	120	125
5	Asp Pro Arg Asp Arg		
	130		
10	<210> 19		
	<211> 133		
	<212> PRT		
	<213> Lycopersicon esculentum		
15	<400> 19		
	Lys His Ile Ile His Arg Asp Leu Lys Pro Ser Asn Leu Leu Ile Asn		
	1	5	10 15
20	His Arg Gly Asp Val Lys Ile Thr Asp Phe Gly Val Ser Ala Val Leu		
	20	25	30
25	Ala Ser Thr Ser Gly Leu Ala Asn Thr Phe Val Gly Thr Tyr Asn Tyr		
	35	40	45
30	Met Ser Pro Glu Arg Ile Ser Gly Gly Ala Tyr Asp Tyr Lys Ser Asp		
	50	55	60
35	Ile Trp Ser Leu Gly Leu Val Leu Leu Glu Cys Ala Thr Gly His Phe		
	65	70	75 80
40	Pro Tyr Lys Pro Pro Glu Gly Asp Glu Gly Trp Val Asn Val Tyr Glu		
	85	90	95
45	Leu Met Glu Thr Ile Val Asp Gln Pro Glu Pro Cys Ala Pro Pro Asp		
	100	105	110
50	Gln Phe Ser Pro Gln Phe Cys Ser Phe Ile Ser Ala Cys Val Gln Lys		
	115	120	125
55	His Gln Lys Asp Arg		
	130		
50	<210> 20		
	<211> 132		
	<212> PRT		
	<213> Zea mays		
55	<400> 20		
	Arg His Val Ile His Arg Asp Ile Lys Pro Ser Asn Leu Leu Val Asn		
	1	5	10 15



EP 1 078 985 A2

5 Lys Lys Gly Glu Val Lys Ile Thr Asp Phe Gly Val Ser Ala Val Leu  
 20 25 30  
 Ala Ser Ser Ile Gly Gln Arg Asp Thr Phe Val Gly Thr Tyr Asn Tyr  
 35 40 45  
 10 Met Ala Pro Glu Arg Ile Ser Gly Ser Thr Tyr Asp Tyr Lys Ser Asp  
 50 55 60  
 15 Ile Trp Ser Leu Gly Leu Val Ile Leu Glu Cys Ala Ile Gly Arg Phe  
 65 70 75 80  
 Pro Tyr Ile Pro Ser Glu Gly Glu Gly Trp Leu Ser Phe Tyr Glu Leu  
 85 90 95  
 20 Leu Glu Ala Ile Val Asp Gln Pro Pro Pro Ser Ala Pro Ala Asp Gln  
 100 105 110  
 25 Phe Ser Pro Glu Phe Cys Ser Phe Ile Ser Ser Cys Ile Gln Lys Asp  
 115 120 125  
 Pro Ala Gln Arg  
 130  
 30  
 <210> 21  
 <211> 88  
 <212> PRT  
 35 <213> Unknown  
 <220>  
 <223> Description of Unknown Organism: another MAPKK  
 40 gene  
 <400> 21  
 Asp Thr Phe Thr Gly Thr Tyr Asn Tyr Met Ala Pro Glu Arg Ile Ser  
 45 1 5 10 15  
 Gly Gln Lys His Gly Tyr Met Ser Asp Ile Trp Ser Leu Gly Leu Val  
 20 25 30  
 50 Met Leu Glu Leu Ala Thr Gly Glu Phe Pro Tyr Pro Pro Arg Glu Ser  
 35 40 45  
 55 Phe Tyr Glu Leu Leu Glu Ala Val Val Asp His Pro Pro Pro Ser Ala  
 50 55 60

EP 1 078 985 A2

Pro Ser Asp Gln Phe Ser Glu Glu Phe Cys Ser Phe Val Ser Ala Cys  
65 70 75 80

Ile Gln Lys Asn Ala Ser Asp Arg  
85

<210> 22

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 22

gtatgtgccg acaaagtcac tggccagtcc atctgtgctt gctagtactg cactcacac 59

<210> 23

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 23

gtactagcaa gcacagatgg actggccaat gactttgtcg gcacatacaa ctatatgtc 59

<210> 24

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:nucleic acid  
sequence

<400> 24

ctctagagga tccccgggtg gtcagtcct t 31

Claims

1. A nucleic acid sequence encoding a derivative of a plant mitogen-activated protein kinase kinase, wherein said

derivative contains a negative charge at a core phosphorylation site of said protein kinase kinase.

2. The nucleic acid sequence of claim 1, wherein said derivative comprises replacement of one or more amino acids with an amino acid selected from the group consisting of: aspartic acid and glutamic acid.

3. The nucleic acid sequence of claim 2, wherein said derivative comprises replacement of one or more serine or threonine amino acids with an amino acid selected from the group consisting of: aspartic acid and glutamic acid.

4. The nucleic acid sequence of claim 3, wherein said nucleic acid sequence is isolated from the group consisting of: *Arabidopsis thaliana*, *Lycopersicum esculentum*, *Zea mais*, *N tabucum*, *D discoideum* and *Leishmania donovani*.

5. The nucleic acid sequence of claim 4, wherein said one or more threonine or serine amino acids are selected from the group consisting of:

*Lycopersicum esculentum* c.v. Bonny Best, tMEK 2: 219serine, 220threonine, 221 serine and 226threonine;  
*Arabidopsis thaliana*, AtMAP2K $\alpha$ : 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMKK4: 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMEK1: 219serine, 220threonine, 221serine, 222serine and 226serine;  
*L. esculentum*, LeMEK1: 219serine, 220threonine, 221serine and 226threonine;  
*Zea mais*, ZmMEK1: 219serine, 220serine and 226threonine;  
*A. thaliana*, At MAP2K $\beta$ : 218threonine, 220threonine and 226threonine;  
*N tabucum*, NPK2: 219serine, 220serine and 226threonine;  
*A. thaliana*, AtMKK3: 220serine and 226threonine;  
*D discoideum*, DdMEK1, 220threonine, 222serine and 226threonine; and  
*Leishmania donovani*, LPK: 220threonine, 224serine, 225serine and 226threonine;

wherein the amino acid numbering system is based on the tomato gene tMEK2.

6. The nucleic acid sequence of claim 5, wherein the nucleic acid is from tomato cv. Bonny Best, and wherein in the encoded derivative amino acids serine221 and threonine226 have been replaced with aspartic acid.

7. A derivative of a plant mitogen-activated protein kinase kinase, wherein said derivative contains a negative charge in a core phosphorylation site of said protein kinase kinase.

8. The derivative of claim 7, wherein said derivative comprises replacement of one or more amino acids with an amino acid selected from the group consisting of: aspartic acid and glutamic acid.

9. The derivative of claim 8, wherein said derivative comprises replacement of one or more serine or threonine amino acids with an amino acid selected from the group consisting of: aspartic acid and glutamic acid.

10. The derivative of claim 9, wherein the mitogen-activated protein kinase kinase is derived from plants selected from the group consisting of: *Arabidopsis thaliana*, *Lycopersicum esculentum*, *Zea mais*, *N tabucum*, *D discoideum* and *Leishmania donovani*.

11. The derivative of claim 10, wherein one or more of said serine or threonine amino acids are selected from the group consisting of:

*Lycopersicum esculentum* c.v. Bonny Best, tMEK 2: 219serine, 220threonine, 221serine and 226threonine;  
*Arabidopsis thaliana*, AtMAP2K $\alpha$ : 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMKK4: 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMEK1: 219serine, 220threonine, 221serine, 222serine and 226serine;  
*L. esculentum*, LeMEK1: 219serine, 220threonine, 221serine and 226threonine;  
*Zea mais*, ZmMEK1: 219serine, 220serine and 226threonine;  
*A. thaliana*, At MAP2K $\beta$ : 218threonine, 220threonine and 226threonine;  
*N tabucum*, NPK2: 219serine, 220serine and 226threonine;  
*A. thaliana*, AtMKK3: 220serine and 226threonine;  
*D discoideum*, DdMEK1, 220threonine, 222serine and 226threonine; and

*Leishmania donovani*, LPK: 220threonine, 224serine, 225serine and 226threonine;

wherein the amino acid numbering system is based on the tomato gene tMEK2.

12. The derivative of claim 11, wherein the derivative of a mitogen-activated protein kinase kinase is derived from tomato cv. Bonny Best, and wherein the amino acids serine221 and threonine226 have been replaced with aspartic acid.
13. A cloning vector comprising the nucleic acid sequence of claim 1.
14. A transgenic plant comprising the cloning vector of claim 13.
15. A transgenic plant comprising the nucleic acid sequence of claim 1.
16. A method of increasing disease resistance or enhancing stress tolerance in a plant by introducing into said plant a nucleic acid sequence encoding a derivative of a mitogen-activated protein kinase kinase, wherein said derivative contains a negative charge at a core phosphorylation site of said protein kinase kinase.
17. The method of claim 16, wherein said derivative comprises replacement of one or more amino acids with an amino acid selected from the group consisting of: aspartic acid and glutamic acid.
18. The method of claim 17, wherein said derivative comprises replacement of one or more serine or threonine amino acids with an amino acid selected from the group consisting of: aspartic acid and glutamic acid.
19. The method of claim 18, wherein said nucleic acid is isolated from the group consisting of: *Arabidopsis thaliana*, *Lycopersicum esculentum*, *Zea mais*, *N tabacum*, *D discoideum*, *Leishmania donovani*, *Drosophila melanogaster*, *Homo sapiens*, *R norvegicus*, *Saccharomyces cerevisiae* and *Candida albicans*.
20. The method of claim 19, wherein said one or more serine or threonine amino acids are selected from the group consisting of:

*Lycopersicum esculentum*, tMEK 2: 219serine, 220threonine, 221serine and 226threonine;  
*Arabidopsis thaliana*, AtMAP2K $\alpha$ : 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMKK4: 220threonine, 226serine and 227serine;  
*A. thaliana*, AtMEK1: 219serine, 220threonine, 221serine, 222serine and 226serine;  
*L. esculentum*, LeMEK1: 219serine, 220threonine, 221serine and 226threonine;  
*Zea mais*, ZmMEK1: 219serine, 220serine and 226threonine;  
*A. thaliana*, At MAP2K $\beta$ : 218threonine, 220threonine and 226threonine;  
*N tabacum*, NPK2: 219serine, 220serine and 226threonine;  
*A. thaliana*, AtMKK3: 220serine and 226threonine;  
*D discoideum*, DdMEK1, 220threonine, 222serine and 226threonine;  
*Leishmania donovani*, LPK: 220threonine, 224serine, 225serine and 226threonine;  
*Drosophila melanogaster*, HEP: 220serine and 226threonine; human, MEK1: 220serine and 226serine;  
*R norvegicus*, MEK5: 220serine and 226threonine;  
*Homo sapiens*, MKK3: 220serine and 226threonine;  
*Saccharomyces cerevisiae*, PBS2: 220serine and 226threonine;  
*S. cerevisiae*, STE7: 220serine and 226threonine;  
*Candida albicans*, HST 7: 220serine and 226threonine; and  
*S. cerevisiae*, MKK1: 220serine, 225threonine and 226threonine;

wherein the amino acid numbering system is based on the tomato gene tMEK2.

21. The method of claim 20, wherein the nucleic acid is from tomato cv. Bonny Best, and wherein in the encoded derivative amino acids serine221 and threonine226 have been replaced with aspartic acid.
22. The method of claim 16, wherein said nucleic acid is introduced by a method selected from transformation and particle bombardment.

1	ATSAAGAAAGGATCTTTTGCACCTAATCTTAACTCTCTCTCTCTCTCTCTCTCTGATGAAGTT	50
1	M K K G S F A P N L K L S L P P P D E V	20
51	GCTCTCTCCAAATCTCTGACTGAATCAGGAACATTTAAGGATGAGATCTTTCTGGTGAAT	120
21	A L S K F L T E S G T F K D G D L L V H	40
121	AGAGATCGAGTTCGAATTGTTTCGCAGAGTGAAGTTCCAGCTCTTCAGTTATACAGCCA	180
41	R D G V R I V S Q S E V A A P S V I Q P	60
191	TCAGACAACCAAGTTATGCTTAGCTGATTTTGAAGCAGTAAAGTTATTGGAAAGGGAAT	240
51	S D N Q L C L A D F E A V K V I G K G H	80
241	GCTGGTATAGTCCGGCTGGTTCAGCATAAATCGACAGGGCAATTTTTCGCTCTCAAGTT	300
91	G G I V R L V Q H K W T G Q F F A L K V	100
301	ATTGAGATGAATATTGATGAGTCTATCGGCAACATATTGCTCAAGAACTGAGAATTAAT	360
101	I Q M N I D E S H R K H I A Q E L R I H	120
361	CAGTCATCCCAAGTGTCCATATGTTGTCATATGCTATCAGTCTCTTCGACAATGCTGT	420
121	Q S S Q C P Y V V I C Y Q S F F D N G A	140
421	ATATCCTTGATTTTGGAGTATATGGATGGTGGTTCCTTAGCAGATTTTCTGAAAAGGTC	480
141	I S L E L E Y M O G G S L A D F L K K V	160
481	AAAAAATACCTGAACGATTCTTCTGCTTATCTGCAACAGGTTCTCAAGGCTTGTGG	540
161	K T I P E R F L A V I C K Q V L R G L W	180
541	TATCTTCATCATGAGAAGCATATTATTCACAGGATTGAAACCTTCGAATTTGCTAATC	600
181	Y L H H E K H I E H R O L K P S N L L I	200
601	AATCAGAGGCTGATGTCAAAATCAGAGCTTTGGTGTGAGTGCAGTACTAGCAAGCACA	660
201	N H R G D V K I T D F G V S A V L A S T	220
661	TCTGGACTGGCCAATACCTTTGTGGCACATACAATATATGCTTCAGAGAGAATTCA	720
221	S G L A N T F V G T Y N Y M S P E R I S	240
721	GGAGGTGCTATGATTACAAAAGCGACATTTGGAGCTTGGGTTTAGTCTTGTCTCGAGTGT	780
241	G G A Y D Y K S D I H S L G L V L L E C	260
781	GCAACAGGTCATTTCCCATATAAACACCCGAGGGAGATGAAGGATGGGTCAATGTCTAT	840
261	A T G H F P Y K P P E G D E G M V N V Y	280
941	GAACTTATGGAAACCATAGTTGACCAACCAGAACCTTGTGCACCTCTGACCAATTTCT	900
281	E L H E T I V D Q P E P C A P P D Q F S	300
901	CCACAATTCTGCTCATTCATATCTGCATGTGTCCAGAAGCACCAGAAGGACAGACTGTCTG	960
301	P Q F C S F I S A C V Q K H Q K D R L S	320
961	GCAATGATCTCATGAGTCAACCTTTTCATCACCATGTACGATGACCAGGATATCGATCTT	1020
321	A N D L H S H P F I T H Y D D Q D I D L	340
1021	GGATCTTACTTCACTTCCGAGGACCTCCATTGGCAACACTTACTGAGCTATAA	1074
341	G S Y F T S A G P P L A T L T E L	358

FIGURE 1a

[illegible]

**FIGURE 1b**

1. tMEK2	214	SAVLA <del>ST</del> SGLAN <del>TF</del>	227	tomato cv Bonny Best
2. AtMAP2K		SRILAQTMDPCNSS		Arabidopsis
3. AtMKK4		SRILAQTMDPCNSS		Arabidopsis
4. AtMEK1		SKILT <del>ST</del> SSLAN <del>SF</del>		Arabidopsis
5. LeMEK1		SAVLA <del>ST</del> SGLAN <del>TF</del>		tomato cv Ailsa Craig
6. ZmMEK1		SAVLASSIGQRDTF		maize
7. AtMAP2K		STVMTNTAGLAN <del>TF</del>		Arabidopsis
8. NPK2		SAGLESSIAMCATF		tobacco
9. AtMKK3		SAGLENSMAMCATF		Arabidopsis
10. DdMEK1		SGQLQH <del>TL</del> SKAV <del>TW</del>		<i>D. discoideum</i>
11. LPK		S-KLIQ <del>TL</del> AVS <del>STY</del>		<i>leishmania donovani</i>
12. HEP		SGRLVDSK-ANTR		<i>Drosophila</i>
13. MEK1		SGQLIDSM-AN <del>SF</del>		human
14. MEK5		STQLVNSI-AK <del>TY</del>		rat
15. MKK3		SGYLVDSV-AK <del>TM</del>		human
16. PBS2		SGNLVASL-AK <del>TN</del>		yeast
17. STE7		SKKLINSI-AD <del>TF</del>		yeast
18. HST7		SREL <del>TN</del> SLAMAD <del>TF</del>		<i>Candida albicans</i>
19. MKK1		SGEAVNSL-AT <del>TF</del>		yeast

FIGURE 1c




FIGURE 2



# Control Construct



## tMEK2<sup>WT</sup> Construct



## tMEK2<sup>MUT</sup> Construct



FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6



FIGURE 7